

Existence Theorems for Elliptic Quasi-Variational Inequalities in Banach Spaces

Risei Kano*, Nobuyuki Kenmochi[†] and Yusuke Murase[#]

^{*,#}Department of Mathematics, Graduate School of Science & Technology
Chiba University, 1-33 Yayoi-cho, Inage-ku, Chiba, 263-8522 Japan

[†]Department of Mathematics, Faculty of Education, Chiba University
1-33 Yayoi-cho, Inage-ku, Chiba, 263-8522 Japan

Abstract. A class of quasi-variational inequalities (QVI) of the elliptic type is studied in Banach spaces. The concept of QVI was earlier introduced by A. Bensoussan and J. L. Lions [2] and its general theory was evolved by many mathematicians, for instance, see [7,9,13] and a monograph [1]. In this paper we give a generalization of the existence theorem due to J. L. Joly and U. Mosco [6,7] from not only the view-point of the nonlinear operator theory, but also the application to nonlinear variational inequalities including partial differential operators. In fact, employing the compactness argument based on the Mosco convergence (cf.[11]) for convex sets and the graph convergence for nonlinear operators, we shall prove an abstract existence result for our class of QVI's. Moreover we shall give some new applications to QVI's arising in the material science.

1. Introduction

Let X be a real reflexive Banach space and X^* be its dual. We assume that X and X^* are strictly convex and denote by $\langle \cdot, \cdot \rangle$ the duality pairing between X^* and X . Given a nonlinear operator A from X into X^* , an element $g^* \in X^*$ and a closed convex subset K of X , the variational inequality is formulated as a problem to find u in X such that

$$u \in K, \quad \langle Au - g^*, u - w \rangle \leq 0, \quad \forall w \in K. \quad (1.1)$$

Variational inequality has been studied by many mathematicians, for instance see J. L. Lions and G. Stampacchia [10], F. Browder [5], H. Brézis [4] and their references.

The concept of quasi-variational inequality was introduced by A. Bensoussan and J. L. Lions [2] in order to solve some problems in the control theory. Given an operator

$A : X \rightarrow X^*$, an element $g^* \in X^*$ and a family $\{K(v); v \in X\}$ of closed convex subsets of X , the quasi-variational inequality is a problem to find u in X such that

$$u \in K(u), \quad \langle Au - g^*, u - w \rangle \leq 0, \quad \forall w \in K(u). \quad (1.2)$$

As is seen from (1.2), the constraint $K(u)$ for the quasi-variational inequality depends upon the unknown u , which causes one of main difficulties in the mathematical treatment of quasi-variational inequalities. The theory of quasi-variational inequality has been evolved for various classes of the mapping $v \rightarrow K(v)$ and the linear or nonlinear operator $A : X \rightarrow X^*$; see for instance [6,7,13], in which two approaches to quasi-variational inequalities were proposed. One of them is the so-called monotonicity method in Banach lattices X (cf. [13]), and for the mapping $v \rightarrow K(v)$ the monotonicity condition

$$\min\{w_1, w_2\} \in K(v_1), \quad \max\{w_1, w_2\} \in K(v_2), \quad \text{if } v_1, v_2 \in X \text{ with } v_1 \leq v_2, \quad (1.3)$$

is required, and an existence result for (1.2) is proved with the help of a fixed point theorem in Banach lattices. Another is the so-called compactness method in which some compactness properties are required for the mapping $v \rightarrow K(v)$ such as $K(v_n)$ converges to $K(v)$ in the Mosco sense, if $v_n \rightarrow v$ weakly in X as $n \rightarrow \infty$. In this framework, an existence result for (1.2) was shown by J. L. Joly and U. Mosco [7]. However this result seems not enough from some points of applications. The objective of this paper is to generalize the result in [7] to the case that $A : X \rightarrow X^*$ is the multivalued pseudo-monotone operator, $Au = \tilde{A}(u, u)$, generated by a semimonotone operator $\tilde{A} : X \times X \rightarrow X^*$. In such a case our quasi-variational inequality is of the form: Find $u \in X$ and $u^* \in X^*$ such that

$$u \in K(u), \quad u^* \in Au, \quad \langle u^* - g^*, u - w \rangle \leq 0, \quad \forall w \in K(u). \quad (1.4)$$

This generalization (1.4) is new and enables us to apply it to the following quasi-variational inequality arising in the elastic-plastic torsion problem for visco-elastic material : Find $u \in H_0^1(\Omega)$ and $\tilde{u} \in L^2(\Omega)$ satisfying

$$\left\{ \begin{array}{l} |\nabla u| \leq k_c(u) \text{ a.e. on } \Omega, \quad \tilde{u} \in \beta(u) \text{ a.e. on } \Omega, \\ \sum_{i,j=1}^N \int_{\Omega} a_{ij}(x, u) \frac{\partial u}{\partial x_i} \frac{\partial(u-w)}{\partial x_j} dx + \int_{\Omega} \tilde{u}(u-w) dx \leq \int_{\Omega} f(u-w) dx, \\ \forall w \in H_0^1(\Omega) \text{ with } |\nabla w| \leq k_c(u) \text{ a.e. on } \Omega, \end{array} \right. \quad (1.5)$$

where Ω is a bounded smooth domain in \mathbf{R}^N , f is given in $L^2(\Omega)$, $k_c(\cdot)$ is a positive, smooth and bounded function on \mathbf{R} and $\beta(\cdot)$ is a maximal monotone graph in $\mathbf{R} \times \mathbf{R}$ with linear growth at $\pm\infty$. In this case our abstract result is applied to

$$K(v) := \{w \in H_0^1(\Omega); |\nabla w| \leq k_c(v) \text{ a.e. on } \Omega\} \quad (1.6)$$

and

$$\tilde{A}(v, u) := - \sum_{i,j=1}^N \frac{\partial}{\partial x_j} \left(a_{ij}(x, v) \frac{\partial u}{\partial x_i} \right) + \beta(u); \quad (1.7)$$

it should be noted that the family $\{K(v); v \in H_0^1(\Omega)\}$ given by (1.6) does not satisfy the monotonicity condition (1.3), and that the term $\beta(u)$ of (1.7) is in general multivalued.

Therefore, (1.5) is a new application in the respect that the differential form in (1.7) is quasi-linear and the additional term is multivalued in general.

2. Main results

Let X be a real Banach space and X^* be its dual space, and assume that X and X^* are strictly convex. We denote by $\langle \cdot, \cdot \rangle$ the duality pairing between X^* and X , and by $|\cdot|_X$ and $|\cdot|_{X^*}$ the norms of X and X^* , respectively. For various general concepts on nonlinear multivalued operators from X into X^* , for instance, monotonicity and maximal monotonicity of operators, we refer to the monograph [1]. In this paper, we mean that operators are multivalued, in general. Given a general nonlinear operator A from X into X^* , we use the notations $D(A)$, $R(A)$ and $G(A)$ to denote its domain, range and graph of A . In this paper, we formulate quasi-variational inequalities for a class of nonlinear operators, which is called semimonotone, from $X \times X$ into X^* .

Definition 2.1. An operator $\tilde{A}(\cdot, \cdot) : X \times X \rightarrow X^*$ is called *semimonotone*, if $D(\tilde{A}) = X \times X$ and the following conditions (SM1) and (SM2) are satisfied:

(SM1) For any fixed $v \in X$ the mapping $u \rightarrow \tilde{A}(v, u)$ is maximal monotone form $D(\tilde{A}(v, \cdot)) = X$ into X^* .

(SM2) Let u be any element of X and $\{v_n\}$ be any sequence in X such that $v_n \rightarrow v$ weakly in X . Then, for every $u^* \in \tilde{A}(v, u)$ there exists a sequence $\{u_n^*\}$ in X^* such that $u_n^* \in \tilde{A}(v_n, u)$ and $u_n^* \rightarrow u^*$ in X^* as $n \rightarrow +\infty$.

Let $\tilde{A} : D(\tilde{A}) := X \times X \rightarrow X^*$ be a semimonotone operator. Then we define $A : D(A) = X \rightarrow X^*$ by putting $Au := \tilde{A}(u, u)$ for all $u \in X$, which is called the operator generated by \tilde{A} .

Now, for an operator A generated by semimonotone operator, any $g^* \in X^*$ and a mapping $v \rightarrow K(v)$ we consider a quasi-variational inequality, denoted by $P(g^*)$, to find $u \in X$ and $u^* \in X^*$ such that

$$P(g^*) \quad u \in K(u), \quad u^* \in Au, \quad \langle u^* - g^*, w - u \rangle \leq 0, \quad \forall w \in K(u). \quad (2.1)$$

Our main results of this paper are stated as follows.

Theorem 2.1. Let $\tilde{A} : D(\tilde{A}) = X \times X \rightarrow X^*$ be a bounded semimonotone operator and A be the operator generated by \tilde{A} . Let K_0 be a bounded, closed and convex set in X . Suppose that to each $v \in K_0$ a non-empty, bounded, closed and convex subset $K(v)$ of K_0 is assigned, and the mapping $v \rightarrow K(v)$ satisfies the following continuity properties (K1) and (K2) :

(K1) If $v_n \in K_0$, $v_n \rightarrow v$ weakly in X (as $n \rightarrow \infty$), then for each $w \in K(v)$ there is a sequence w_n in X such that $w_n \in K(v_n)$ and $w_n \rightarrow w$ (strongly) in X .

(K2) If $v_n \rightarrow v$ weakly in X , $w_n \in K(v_n)$ and $w_n \rightarrow w$ weakly in X , then $w \in K(v)$.

Then, for any $g^* \in X^*$, the quasi-variational inequality $P(g^*)$ has at least one solution u .

The following theorem is a slightly general version of Theorem 2.1.

Theorem 2.2. Let $\tilde{A} : D(\tilde{A}) = X \times X \rightarrow X^*$ be a bounded semimonotone operator and A be the operator generated by \tilde{A} . Suppose that to each $v \in X$ a non-empty, bounded, closed and convex subset $K(v)$ of X is assigned and there is a bounded, closed and convex subset G_0 of X such that

$$K(v) \cap G_0 \neq \emptyset, \quad \forall v \in X, \quad (2.2)$$

and

$$\inf_{w^* \in Aw} \frac{\langle w^*, w - v \rangle}{|w|_X} \rightarrow \infty \quad \text{as } |w|_X \rightarrow \infty \quad \text{uniformly in } v \in G_0. \quad (2.3)$$

Furthermore, the mapping $v \rightarrow K(v)$ satisfies the following condition (K'1) and the same condition (K2) as in Theorem 2.1.:

(K'1) If $v_n \rightarrow v$ weakly in X , then for each $w \in K(v)$ there is a sequence w_n in X such that $w_n \in K(v_n)$ and $w_n \rightarrow w$ in X .

Then, for any $g^* \in X^*$, the quasi-variational inequality $P(g^*)$ has at least one solution u .

In our proof of Theorems 2.1 and 2.2 we use some results on nonlinear operators of monotone type, which are mentioned below.

Proposition 2.1. Let $\tilde{A} : D(\tilde{A}) = X \times X \rightarrow X^*$ be a semimonotone operator and let $A : X \rightarrow X^*$. Then, the following two properties (a) and (b) are satisfied:

(a) For any $v, u \in X$, $A(v, u)$ is a non-empty, closed, bounded and convex subset of X^* .

(b) Let $\{u_n\}$ and $\{v_n\}$ be sequences in X such that $u_n \rightarrow u$ weakly in X and $v_n \rightarrow v$ weakly in X (as $n \rightarrow \infty$). If $u_n^* \in \tilde{A}(v_n, u_n)$, $u_n^* \rightarrow g$ weakly in X^* and $\limsup_{n \rightarrow \infty} \langle u_n^*, u_n \rangle \leq \langle g, u \rangle$, then $g \in A(v, u)$ and $\lim_{n \rightarrow \infty} \langle u_n^*, u_n \rangle = \langle g, u \rangle$.

Proof. Property (a) immediately follows from the maximal monotonicity of $\tilde{A}(v, \cdot)$ for each $v \in X$. Now we show (b). Assume that $\{u_n\}$, $\{v_n\}$ and $\{u_n^*\}$ are such as in the statement of (b), namely,

$$u_n \rightarrow u \text{ weakly in } X, \quad v_n \rightarrow v \text{ weakly in } X, \quad u_n^* \rightarrow g \text{ weakly in } X^* \quad (2.4)$$

and

$$u_n^* \in \tilde{A}(v_n, u_n), \quad \limsup_{n \rightarrow \infty} \langle u_n^*, u_n \rangle \leq \langle g, u \rangle. \quad (2.5)$$

Now we note from (SM1) that

$$\langle u_n^* - w_n^*, u_n - w \rangle \geq 0, \quad \forall w \in X, \quad \forall w_n^* \in \tilde{A}(v_n, w). \quad (2.6)$$

For any $w \in X$ and any $w^* \in \tilde{A}(v, w)$, use (SM2) to choose a sequence $\{\tilde{w}_n^*\}$ with $\tilde{w}_n^* \in \tilde{A}(v_n, w)$ and $\tilde{w}_n^* \rightarrow w^*$ in X^* . Then, by substituting this sequence w_n^* and letting $n \rightarrow \infty$ in (2.6) we have with the help of (2.4) and (2.5)

$$\langle g - w^*, u - w \rangle \geq 0, \quad \forall w \in X, \quad \forall w^* \in \tilde{A}(v, w).$$

This implies that $g \in \tilde{A}(v, u)$, since $\tilde{A}(v, \cdot)$ is maximal monotone. Corresponding to this $g \in \tilde{A}(v, u)$, by (SM2) choose a sequence $g_n \in \tilde{A}(v_n, u)$ such that $g_n \rightarrow g$ in X^* . Then, by taking $w = u$ and $w_n^* = g_n$ and letting $n \rightarrow \infty$ in (2.6) we obtain

$$\liminf_{n \rightarrow \infty} \langle u_n^*, u_n - u \rangle \geq \liminf_{n \rightarrow \infty} \langle g_n, u_n - u \rangle = 0.$$

namely, $\liminf_{n \rightarrow \infty} \langle u_n^*, u_n \rangle \geq \langle g, u \rangle$. Therefore, on account of (2.5), it holds that

$$\lim_{n \rightarrow \infty} \langle u_n^*, u_n \rangle = \langle g, u \rangle.$$

Thus (b) has been seen. \diamond

We consider a class of nonlinear operators $A : D(A) = X \rightarrow X^*$ satisfying the following properties (PM1) and (PM2):

(PM1) For any $u \in X$, Au is a non-empty, closed, bounded and convex subset of X^* .

(PM2) Let $\{u_n\}$ be a sequence in X such that $u_n \rightarrow u$ weakly in X . If $u_n^* \in Au_n$, $u_n^* \rightarrow g$ weakly in X^* and $\limsup_{n \rightarrow \infty} \langle u_n^*, u_n \rangle \leq \langle g, u \rangle$, then $g \in Au$ and $\lim_{n \rightarrow \infty} \langle u_n^*, u_n \rangle = \langle g, u \rangle$.

This class of nonlinear operators A is called *pseudo-monotone*. The above proposition says that the operator generated by semimonotone \tilde{A} is pseudo-monotone. As to pseudo-monotone operators we refer to [4,8] for fundamental results on their ranges.

Proposition 2.2. *Let $A_1 : D(A_1) \subset X \rightarrow X^*$ be a maximal monotone operator and $A_2 : D(A_2) = X \rightarrow X^*$ be a maximal monotone operator. Suppose that*

$$\inf_{v_1^* \in A_1 v, v_2^* \in A_2 v} \frac{\langle v_1^* + v_2^*, v - v_0 \rangle}{|v|_X} \rightarrow \infty \text{ as } |v|_X \rightarrow \infty, v_1 \in D(A_1).$$

Then $R(A_1 + A_2) = X^$.*

For a proof of Proposition 2.2, see [4,5,8].

3. Proof of main theorems

We begin with the proof of Theorem 2.1.

Proof of Theorem 2.1: The theorem is proved in the following two steps: (A) The case when $\tilde{A}(v, \cdot)$ is strictly monotone from X into X^* for every $v \in X$; (B) The general case as in the statement of Theorem 2.1.

(In the case of (A))

Let v be any element in K_0 and fix it. We consider the variational inequality with state constraint $K(v)$, namely, to find $u \in X$ and $u^* \in X^*$ such that

$$u \in K(v), \quad u^*(v) \in \tilde{A}(v, u), \quad \langle u^*(v) - g^*, u - v \rangle \leq 0, \quad \forall w \in K(v). \quad (3.1)$$

This problem is written in the following form equivalent to (3.1):

$$g^* \in \tilde{A}(v, u) + \partial I_{K(v)}(u), \quad (3.2)$$

where $\partial I_{K(v)}(\cdot) : D(\partial I_{K(v)}) \rightarrow X^*$ is the subdifferential of the indicator function of $K(v)$, i.e.

$$I_{K(v)}(z) := \begin{cases} 0 & \text{if } z \in K(v), \\ \infty & \text{if } z \in X - K(v); \end{cases}$$

note that $\partial I_{K(v)}$ is maximal monotone. It follows from Proposition 2.2 that $R(\tilde{A}(v, \cdot) + \partial I_{K(v)}) = X^*$; in fact, the coerciveness condition of Proposition 2.2 for $A_1 := \partial I_{K(v)}$ and $A_2 := A(v, \cdot)$ is automatically satisfied, since $D(A_1) = K(v)$ is bounded in X . Hence there exists an element u which satisfies (3.2) (therefore (3.1)) for each $v \in K_0$. Moreover, the solution u is unique by the strict monotonicity of $\tilde{A}(v, \cdot)$ and $u \in K_0$. Using this fact, we define a mapping S from K_0 into itself which assigns to each $v \in K_0$ the solution $u \in K_0$ of (3.1), i.e. $u = Sv$.

Next, we show that S is weakly continuous in K_0 . Let $\{v_n\}$ be any sequence in K_0 such that $v_n \rightarrow v$ weakly in X , and put $u_n = Sv_n$ ($\in K_0$) for $n = 1, 2, \dots$. Now, let $\{u_{n_k}\}$ be any weakly convergent subsequence of $\{u_n\}$ and denote by u the weak limit; note by condition (K2) that $u \in K(v)$. We are going to check that u is a unique solution of (3.1). To do so, first observe that there is $u_n^* \in \tilde{A}(v_n, u_n)$ such that

$$\langle u_n^* - g^*, u_n - w \rangle \leq 0, \quad \forall w \in K(v_n) \quad (3.3)$$

Using condition (K1), we find a sequence $\{\tilde{u}_k\}$ such that $\tilde{u}_k \in K(v_{n_k})$ and $\tilde{u}_k \rightarrow u$ in X (as $k \rightarrow \infty$). By the boundedness of $\tilde{A}(\cdot, \cdot)$, we may assume that $u_{n_k}^* \rightarrow u^*$ weakly in X^* for some $u^* \in X^*$. Now, taking $n = n_k$ and $w = \tilde{u}_k$ in (3.3), we see that

$$\begin{aligned} \limsup_{k \rightarrow \infty} \langle u_{n_k}^*, u_{n_k} \rangle &= \limsup_{k \rightarrow \infty} \{ \langle u_{n_k}^*, u_{n_k} - \tilde{u}_k \rangle + \langle u_{n_k}^*, \tilde{u}_k \rangle \} \\ &\leq \limsup_{k \rightarrow \infty} \{ \langle g^*, u_{n_k} - \tilde{u}_k \rangle + \langle u_{n_k}^*, \tilde{u}_k \rangle \} \\ &= \langle u^*, u \rangle \end{aligned}$$

Therefore it follows from Proposition 2.1 that

$$u^* \in \tilde{A}(v, u), \quad \lim_{k \rightarrow \infty} \langle u_{n_k}^*, u_{n_k} \rangle = \langle u^*, u \rangle. \quad (3.4)$$

We go back to (3.3) with $n = n_k$. For any $w \in K(v)$, we use (K1) to choose a sequence $w_k \in K(v_{n_k})$ such that $w_k \rightarrow w$ in X . Taking $n = n_k$ and $w = w_k$ in (3.3) and passing to the limit as $k \rightarrow \infty$ in (3.3), by (3.4) we obtain the variational inequality (3.1). Thus $u = Sv$, and S is weakly continuous in K_0 .

Since K_0 is a weakly compact and convex set in X , we infer from the well-known fixed-point theorem for compact mappings that S has at least one fixed point in K_0 . This fixed point u is clearly a solution of our quasi-variational inequality $P(g^*)$. \diamond

(In the case of (B))

We approximate $\tilde{A}(v, u)$ by $\tilde{A}_\varepsilon(v, u) := \tilde{A}(v, u) + \varepsilon J(u)$ for any $u, v \in X$ and with parameter $\varepsilon \in (0, 1]$; note that the duality mapping J from X into X^* is strictly monotone and hence \tilde{A}_ε is a semimonotone operator such that $\tilde{A}_\varepsilon(v, \cdot)$ is strictly monotone for every $v \in X$. By the result of the case (A), for each $g^* \in X^*$ there exists a solution $u_\varepsilon \in K_0$ of the quasi-variational inequality

$$u_\varepsilon \in K(u_\varepsilon), \quad u_\varepsilon^* \in Au_\varepsilon, \quad \langle u_\varepsilon^* + \varepsilon Ju_\varepsilon - g^*, u_\varepsilon - w \rangle \leq 0, \quad \forall w \in K(u_\varepsilon), \quad (3.5)$$

where A is the operator generated by \tilde{A} . Now, choose a sequence $\{\varepsilon_n\}$, with $\varepsilon_n \downarrow 0$, such that $u_n := u_{\varepsilon_n} \rightarrow u$ weakly in X for some $u \in K_0$. Then, by conditions (K1) and (K2), we see that $u \in K(u)$ and there is a sequence $\{\tilde{u}_n\}$ such that $\tilde{u}_n \in K(u_n)$ and $\tilde{u}_n \rightarrow u$ in X . Moreover, by the boundedness of $\{u_n^* := u_{\varepsilon_n}^*\}$ in X^* , we may assume that $u_n^* \rightarrow u^*$ weakly in X^* for some $u^* \in X^*$. Substitute u_n and \tilde{u}_n for u_ε and w in (3.5) with $\varepsilon = \varepsilon_n$, respectively, and pass to the limit as $n \rightarrow \infty$ to get

$$\begin{aligned} & \limsup_{n \rightarrow \infty} \langle u_n^*, u_n - u \rangle \\ &= \limsup_{n \rightarrow \infty} \{ \langle u_n^* + \varepsilon_n Ju_n, u_n - \tilde{u}_n \rangle + \langle u_n^* + \varepsilon_n Ju_n, \tilde{u}_n - u \rangle \} \\ &= \limsup_{n \rightarrow \infty} \{ \langle g^*, u_n - \tilde{u}_n \rangle + \langle u_n^*, \tilde{u}_n - u \rangle \} \\ &\leq 0. \end{aligned}$$

Since A is pseudo-monotone from X into X^* (cf. Proposition 2.1), it follows from the above inequality that

$$u^* \in Au, \quad \lim_{n \rightarrow \infty} \langle u_n^*, u_n \rangle = \langle u^*, u \rangle. \quad (3.6)$$

Now, for each $w \in K(u)$, by (K1) we choose $\{\tilde{w}_n\}$ such that $\tilde{w}_n \in K(u_n)$ and $\tilde{w}_n \rightarrow w$ in X , and then substitute them for w in (3.5) with $\varepsilon = \varepsilon_n$ to have

$$\langle u_n^* + \varepsilon_n Ju_n - g^*, u_n - \tilde{w}_n \rangle \leq 0. \quad (3.7)$$

By (3.6), letting $n \rightarrow +\infty$ in (3.7) yields that $\langle u^* - g^*, u - w \rangle \leq 0$. Thus u is a solution of our quasi-variational inequality $P(g^*)$. \diamond

Next we proceed to the proof of Theorem 2.2.

Proof of Theorem 2.2: Put $d_1 := \sup_{w \in G_0} |w|_X$ and

$$d_2 := \sup \left\{ |w|_X; w \in X, \inf_{w^* \in Aw} \frac{\langle w^*, w - v \rangle}{|w|_X} \leq |g^*|_{X^*} (1 + d_1), \forall v \in G_0 \right\}.$$

By condition (2.3), d_2 is finite. Also we put $M_0 := d_1 + d_2 + 1$, and for any number $M \geq M_0$ consider the closed ball $B_M := \{w \in X; |w|_X \leq M\}$ as well as bounded closed and convex sets $K_M(v) := K(v) \cap B_M$ for all $v \in X$. Since $G_0 \subset B_M$, (2.2) implies that $K_M(v)$ is non-empty for every $v \in X$.

We now show that conditions (K1) and (K2) in Theorem 2.1 with $K_0 = B_M$ and $K(\cdot) = K_M(\cdot)$ are satisfied. The verification of (K2) is easy. We check condition (K1)

for $K_0 = B_M$ and $K(\cdot) = K_M(\cdot)$ as follows. Let w be any element in $K_M(v)$. Then, by condition (K'1) for $K(\cdot)$, for a sequence $\{v_n\} \subset B_M$ weakly converging to v there is a sequence $\{w_n\}$ such that $w_n \in K(v_n)$ and $w_n \rightarrow w$ in X . In the case of $|w|_X < M$, we see that $|w_n|_X < M$ and hence $w_n \in K_M(v_n)$ for all large n . In the case of $|w|_X = M$, choose an element $v_0 \in K(v) \cap G_0$ and put

$$w^m := \left(1 - \frac{1}{m}\right)w + \frac{1}{m}v_0, \quad m = 1, 2, \dots \quad (3.8)$$

Clearly $w^m \in K_M(v)$ and $|w^m|_X < M$. Therefore, according to the above argument, for each m there is a sequence $\{w_n^m\}_{n=1}^\infty$ such that $w_n^m \in K_M(v_n)$ and $w_n^m \rightarrow w^m$ in X as $n \rightarrow \infty$. For each m choose a number $n(m)$ so that $|w^m - w_n^m|_X \leq \frac{1}{m}$ for all $n \geq n(m)$. We may choose $\{n(m)\}_{m=1}^\infty$ so that $n(m-1) < n(m)$ for all $m = 0, 1, \dots$, where $n(0) = 1$. We put

$$w_n = w_n^m \quad \text{if} \quad n(m) \leq n < n(m+1), \quad m = 0, 1, \dots \quad (3.9)$$

It is easy to see that $w_n \in K_M(v_n)$ and $w_n \rightarrow w$ in X .

By the above observation we can apply Theorem 2.1 to find an element u_M such that

$$u_M \in K_M(u_M), \quad u_M^* \in Au_M, \quad \langle u_M^* - g^*, u_M - w \rangle \leq 0, \quad \forall w \in K_M(u_M). \quad (3.10)$$

Also, by condition (2.3), $\{u_M; M \geq M_0\}$ is bounded in X , so that there are a sequence $\{M_n\}$ with $M_n \uparrow \infty$ and elements $u \in X$, $u^* \in X^*$ such that $u_n := u_{M_n} \rightarrow u$ weakly in X and $u_n^* := u_{M_n}^* \rightarrow u^*$ weakly in X^* as $n \rightarrow \infty$. We note $u \in K(u)$ by (K2). It follows from (K'1) that for each $w \in K(u)$ there is a sequence $\{\tilde{w}_n\}$ such that $\tilde{w}_n \in K(u_n)$ and $\tilde{w}_n \rightarrow w$ in X . In particular, denote by $\{\tilde{u}_n\}$ the sequence $\{\tilde{w}_n\}$ corresponding to $w = u$. Here, we substitute M_n and \tilde{u}_n for M and w in (3.10) to obtain $\langle u_n^* - g^*, u_n - \tilde{u}_n \rangle \leq 0$. Hence it follows that

$$\begin{aligned} & \limsup_{n \rightarrow \infty} \langle u_n^*, u_n - u \rangle \\ &= \limsup_{n \rightarrow \infty} \{ \langle u_n^* - g^*, u_n - \tilde{u}_n \rangle + \langle u_n^*, \tilde{u}_n - u \rangle + \langle g^*, u_n - \tilde{u}_n \rangle \} \\ &\leq 0. \end{aligned}$$

By the pseudo-monotonicity of A this implies that

$$u^* \in Au, \quad \lim_{n \rightarrow +\infty} \langle u_n^*, u_n \rangle = \langle u^*, u \rangle.$$

By making use of these properties with (K1) and passing to the limit as $n \rightarrow \infty$ in (3.10) with $M = M_n$ and $w = \tilde{w}_n$ as above, we see that $u \in K(u)$ and $\langle u^* - g^*, u - w \rangle \leq 0$ for all $w \in K(u)$. Thus u is a solution of our problem $P(g^*)$. \diamond

4. Application to obstacle problems

In this section, let Ω be a bounded domain in \mathbf{R}^N , $1 \leq N < \infty$, with smooth boundary $\Gamma := \partial\Omega$, and put $X := W_0^{1,p}(\Omega)$ or $W^{1,p}(\Omega)$, $1 < p < \infty$. Let $a_i(x, \eta, \xi)$, $i = 1, 2, \dots, N$, be functions on $\Omega \times \mathbf{R} \times \mathbf{R}^N$ such that

(a1) for all $(\eta, \xi) \in \mathbf{R} \times \mathbf{R}^N$ the function $x \rightarrow a_i(x, \eta, \xi)$ is measurable on Ω for each $i = 1, 2, \dots, N$;

(a2) for a.e. $x \in \Omega$ the function $(\eta, \xi) \rightarrow a_i(x, \eta, \xi)$ is continuous on $\mathbf{R} \times \mathbf{R}^N$ for each $i = 1, 2, \dots, N$;

(a3) there are positive constants c_0, c'_0 and c_1, c'_1 such that

$$c_0|\xi|^{p-1} - c'_0 \leq a_i(x, \eta, \xi) \leq c_1|\xi|^{p-1} + c'_1, \quad i = 1, 2, \dots, N, \quad (4.1)$$

$$\text{a.e. } x \in \Omega, \quad \forall \eta \in \mathbf{R}, \quad \forall \xi = (\xi_1, \xi_2, \dots, \xi_N) \in \mathbf{R}^N;$$

(a4) the following monotonicity property is satisfied:

$$\sum_{i=1}^N \left(a_i(x, \eta, \xi) - a_i(x, \eta, \bar{\xi}) \right) (\xi_i - \bar{\xi}_i) \geq 0, \quad (4.2)$$

$$\text{a.e. } x \in \Omega, \quad \forall \eta \in \mathbf{R}, \quad \forall \xi = (\xi_1, \xi_2, \dots, \xi_N), \quad \forall \bar{\xi} = (\bar{\xi}_1, \bar{\xi}_2, \dots, \bar{\xi}_N) \in \mathbf{R}^N.$$

We define a mapping $\tilde{A}_0(\cdot, \cdot) : X \times X \rightarrow X^*$ by putting

$$\langle \tilde{A}_0(v, u), w \rangle = \sum_{i=1}^N \int_{\Omega} a_i(x, v, \nabla u) \frac{\partial w}{\partial x_i} dx, \quad \forall u, v, w \in X, \quad (4.3)$$

and $A_0 u$ by $\tilde{A}_0(u, u)$ for every $u \in X$. Also, let β be a maximal monotone operator from $D(\beta) = \mathbf{R}$ into \mathbf{R} such that the primitive $\hat{\beta}(r) := \int_0^r \beta(s) ds$ of β satisfies that

$$c_2|r|^p - c'_2 \leq \hat{\beta}(r) \leq c_3|r|^p + c'_3, \quad \forall r \in \mathbf{R}, \quad (4.4)$$

where c_2, c'_2, c_3 and c'_3 are positive constants. Now, we consider an operator $\tilde{A} : X \times X \rightarrow X^*$ given by $\tilde{A}(v, u) := \tilde{A}_0(v, u) + \beta(u)$ for all $v, u \in X$. It is easy to see from (4.3) and (4.4) that \tilde{A} is a bounded and semimonotone operator from $X \times X$ into X^* .

Application 4.1. (Gradient obstacle problem)

Let $X = W_0^{1,p}(\Omega)$, and k_c be a Lipschitz continuous real function on \mathbf{R} such that

$$0 < k_c(r) \leq k^*, \quad \forall r \in \mathbf{R}, \quad (4.5)$$

where k^* is a positive constant, and put

$$K(v) := \{w \in X; |\nabla w| \leq k_c(v) \text{ a.e. on } \Omega\}, \quad \forall v \in X. \quad (4.6)$$

Also, we set

$$K_0 := \{w \in X; |\nabla w| \leq k^* \text{ a.e. on } \Omega\}; \quad (4.7)$$

note from the Sobolev imbedding theorem that K_0 is compact in $C(\bar{\Omega})$.

Lemma 4.1. *The family $\{K(v); v \in X\}$ and the set K_0 , which are respectively given by (4.6) and (4.7), satisfy conditions (K1) and (K2).*

Proof. We prove (K2). Suppose that $v_n \in K_0$, $w_n \in K(v_n)$, $v_n \rightarrow v$ weakly in X and $w_n \rightarrow w$ weakly in X . Then, $v_n \rightarrow v$ in $C(\bar{\Omega})$ and hence $k_c(v_n) \rightarrow k_c(v)$ in $C(\bar{\Omega})$. Therefore, given $\varepsilon > 0$, there exists a positive integer n_ε such that

$$|k_c(v_n) - k_c(v)| \leq \varepsilon \quad \text{on } \Omega, \quad \forall n \geq n_\varepsilon. \quad (4.8)$$

This shows that

$$|\nabla w_n| \leq k_c(v_n) \leq k_c(v) + \varepsilon \quad \text{a.e. on } \Omega, \quad \forall n \geq n_\varepsilon. \quad (4.9)$$

Clearly the set $K_\varepsilon(v) := \{w \in X; |\nabla w| \leq k_c(v) + \varepsilon \text{ a.e. on } \Omega\}$ is bounded, closed and convex in X , so that $K_\varepsilon(v)$ is weakly compact in X . Now, we derive by letting $n \rightarrow +\infty$ in (4.9) that $w \in K_\varepsilon(v)$. Since $\varepsilon > 0$ is arbitrary, we have $w \in K(v)$. Thus (K2) holds.

Next we show (K1). Suppose that $v \in K_0$, $w \in K(v)$ and $\{v_n\} \subset K_0$ such that $v_n \rightarrow v$ weakly in X . By the compactness of K_0 in $C(\bar{\Omega})$ we have that $v_n \rightarrow v$ in $C(\bar{\Omega})$. Since $cw \in K(v)$ for all constant $c \in (0, 1)$ and $cw \rightarrow w$ as $c \uparrow 1$ in X , it is enough to show the existence of a sequence $\{\tilde{w}_n\}$ such that $\tilde{w}_n \in K(v_n)$ and $\tilde{w}_n \rightarrow \tilde{w}$ in X , when $\tilde{w} = cw$ for any $c \in (0, 1)$. In such a case, by conditions (4.5) and (4.8), we can take a small $\varepsilon > 0$ so that $|\nabla \tilde{w}| \leq k_c(v) - \varepsilon$ a.e. on $\bar{\Omega}$. Furthermore, for this $\varepsilon > 0$ we can find a positive integer n_ε such that $k_c(v) \leq k_c(v_n) + \varepsilon$ for all $n \geq n_\varepsilon$. This implies that $|\nabla \tilde{w}| \leq k_c(v_n)$ a.e. on Ω , namely $\tilde{w} \in K(v_n)$ for all $n \geq n_\varepsilon$. Now we define $\{\tilde{w}_n\}$ by putting

$$\tilde{w}_n = \begin{cases} \tilde{w} & \text{for } n \geq n_\varepsilon, \\ \text{some function in } K(v_n) & \text{for } 1 \leq n < n_\varepsilon. \end{cases}$$

Clearly this is a required sequence in condition (K1). \diamond

According to Lemma 4.1, we can apply Theorem 2.1 to solve the following quasi-variational inequality:

$$\begin{cases} u \in X, \quad |\nabla u| \leq k_c(u) \text{ a.e. on } \Omega, \quad u^* \in L^q(\Omega) \text{ with } u^* \in \beta(u) \text{ a.e. on } \Omega; \\ \sum_{i=1}^N \int_{\Omega} a_i(x, u, \nabla u)(u_{x_i} - w_{x_i}) dx + \int_{\Omega} u^*(u - w) dx \leq \int_{\Omega} f(u - w) dx, \\ \forall w \in X \text{ with } |\nabla w| \leq k_c(u) \text{ a.e. on } \Omega, \end{cases} \quad (4.10)$$

where f is given in $L^q(\Omega)$, $\frac{1}{p} + \frac{1}{q} = 1$.

Application 4.2. (Interior obstacle problem)

Let Ω be a one-dimensional bounded open interval, say $(0, 1)$, $X := W^{1,p}(0, 1)$, $1 < p < \infty$, and let $a_1(x, \eta, \xi)$ be a function which satisfy (a1)-(a4) with $N = 1$. Let β be the same as above.

Also, let $k_c(\cdot)$ be a Lipschitz continuous real function on \mathbf{R} such that

$$k_c(r) \leq k^*, \quad \forall r \in \mathbf{R}, \quad (4.11)$$

where k^* is a constant, and put

$$K(v) := \{w \in X; w \geq k_c(v) \text{ on } (0, 1)\}, \quad \forall v \in X. \quad (4.12)$$

Clearly, the constant function k^* belongs to $K(v)$ for all $v \in X$. Futhermore, it follows from (4.1) and (4.4) that

$$\int_0^1 \{a_1(x, v, v_x)v_x + v^*(v - k^*)\} dx \geq c_4|v|_X^p - c'_4, \quad (4.13)$$

$$\forall v \in X, \quad \forall v^* \in L^q(0, 1) \text{ with } v^* \in \beta(v) \text{ a.e. on } (0, 1),$$

where c_4 and c'_4 are positive constants.

Lemma 4.2. *The family $\{K(v); v \in X\}$ and the set $G_0 := \{k^*\}$ satisfy conditions (2.2), (2.3), (K'1) and (K2).*

Proof. We observe that X is compactly embedded in $C([0, 1])$. On account of this fact, the verification of (K'1) and (K2) can be done in a way similar to that in the proof of Lemma 4.1. Also, (2.2) and (2.3) are immediately seen from (4.11), (4.12) and (4.13). \diamond

Now, given $f \in L^q(0, 1)$, by applying Theorem 2.2 we find functions u and u^* such that

$$\left\{ \begin{array}{l} u \in X, \quad u \geq k_c(u) \text{ on } (0, 1), \quad u^* \in L^q(0, 1) \text{ with } u^* \in \beta(u) \text{ a.e. on } (0, 1), \\ \int_0^1 \{a_1(x, u, u_x)(u_x - w_x) + u^*(u - w)\} dx \leq \int_0^1 f(u - w)dx, \\ \forall w \in X \text{ with } w \geq k_c(u) \text{ on } (0, 1). \end{array} \right. \quad (4.14)$$

Application 4.3. (Boundary obstacle problem)

We consider a quasi-variational inequality with constraints on the boundary. Let $X := W^{1,p}(0, 1)$, $1 < p < \infty$, and let $a_1(x, \eta, \xi)$ be a function satisfying condition (a1)-(a4) with $N = 1$. Let β be the same as above. Also, let $k_c^i(\cdot)$, $i = 0, 1$, be two Lipschitz continuous real functions on \mathbf{R} such that

$$k_c^i(r) \leq k^*, \quad \forall r \in \mathbf{R}, \quad i = 0, 1,$$

where k^* is a constant. We define

$$K(v) := \{w \in X; w(i) \geq k_c^i(v(i)), \quad i = 0, 1\}, \quad \forall v \in X, \quad (4.14)$$

and $G_0 = \{k^*\}$, being the singleton set of constant function k^* . Then it is easy to see that G_0 and the family $\{K(v)\}$ satisfy conditions (2.2), (2.3), (K'1) and (K2) in the statement of Theorem 2.2. Therefore, by Theorem 2.2, for each $f \in L^q(0, 1)$, $\frac{1}{p} + \frac{1}{q} = 1$, we find functions u and u^* such that

$$\left\{ \begin{array}{l} u \in X, \quad u(i) \geq k_c(u(i)), \quad i = 0, 1, \quad u^* \in L^q(0, 1) \text{ with } u^* \in \beta(u) \text{ a.e. on } (0, 1); \\ \int_0^1 \{a_1(x, u, u_x)(u_x - w_x) + u^*(u - w)\} dx \leq \int_0^1 f(u - w)dx, \\ \forall w \in X \text{ with } w(i) \geq k_c(u(i)), \quad i = 0, 1. \end{array} \right.$$

In Applications 4.2 and 4.3 we supposed that the space dimension of Ω is one, since we do not know whether condition (K1) or (K'1) holds or not in higher space dimensional cases.

Application 4.4. Next we consider a system of quasi-variational inequality including quasi-linear partial differential operators. Let $a_k(u_1, u_2)$, $k = 1, 2$, be continuous functions with respect to u_1 and u_2 on $\mathbf{R} \times \mathbf{R}$ such that

$$c_5 \leq a_k(u_1, u_2) \leq c'_5, \quad \forall u_1, u_2 \in \mathbf{R},$$

and let $\beta_k(u_k)$, $k = 1, 2$, be maximal monotone operators from $D(\beta_k) = \mathbf{R}$ into \mathbf{R} and the primitives $\hat{\beta}_k(r) := \int_0^r \beta_k(s) ds$ satisfy that

$$c_6|r|^{p_k} - c'_6 \leq \hat{\beta}_k(r) \leq c_7|r|^{p_k} + c'_7, \quad \forall r \in \mathbf{R}, \quad k = 1, 2,$$

where p_k , $k = 1, 2$, are constants satisfying $2 \leq p_k < \infty$. Our problem is formulated in the space $X := W^{1,p_1}(0, 1) \times W^{1,p_2}(0, 1)$, and define an operator $\tilde{A}_0 : X \rightarrow X^*$ by

$$\langle \tilde{A}_0(v, u), w \rangle = \int_0^1 a_1(v_1, v_2) |u_{1,x}|^{p_1-2} u_{1,x} w_{1,x} dx + \int_0^1 a_2(v_1, v_2) |u_{2,x}|^{p_2-2} u_{2,x} w_{2,x} dx,$$

$$\forall v := (v_1, v_2), \quad u := (u_1, u_2), \quad w := (w_1, w_2) \in X.$$

as well as an operator $\tilde{A} : X \rightarrow X^*$ by $\tilde{A}(v, u) := \hat{A}_0(v, u) + (\beta_1(u_1), \beta_2(u_2))$, namely, the first and second component of $\tilde{A}(v, u)$ are respectively written in the form

$$-(a_1(v_1, v_2) |u_{1,x}|^{p_1-2} u_{1,x})_x + \beta_1(u_1) \quad \text{and} \quad -(a_2(v_1, v_2) |u_{2,x}|^{p_2-2} u_{2,x})_x + \beta_2(u_2),$$

which are multivalued, in general, because of $\beta_1(u_1)$ and $\beta_2(u_2)$. As was already seen just before Application 4.1, \tilde{A} is bounded and semimonotone from X into X^* . Further let $k_{kc}(u_1, u_2)$, $k = 1, 2$, be Lipschitz continuous functions with respect to $u = (u_1, u_2)$ on $\mathbf{R} \times \mathbf{R}$ such that $k_{kc} \leq k^*$ on $\mathbf{R} \times \mathbf{R}$ for a constant k^* . Then, given $f_k \in L^{q_k}(0, 1)$, $\frac{1}{p_k} + \frac{1}{q_k} = 1$, $k = 1, 2$, by virtue of Theorem 2.2 there exist vector functions $u = (u_1, u_2) \in X$ and $u^* = (u_1^*, u_2^*) \in L^{q_1}(0, 1) \times L^{q_2}(0, 1)$ such that

$$\left\{ \begin{array}{l} u_k \geq k_{kc}(u_1, u_2) \text{ on } (0, 1), \quad u_k^* \in \beta_k(u_k) \text{ a.e. on } (0, 1), \quad k = 1, 2, \\ \int_0^1 a_k(u_1, u_2) |u_{k,x}|^{p_k-2} (u_{k,x} - w_{k,x}) dx + \int_0^1 u_k^* (u_k - w_k) dx \leq \int_0^1 f_k (u_k - w_k) dx, \\ \forall w_k \in W^{1,p_k}(0, 1) \text{ with } w_k \geq k_{kc}(u_1, u_2) \text{ on } (0, 1), \quad k = 1, 2. \end{array} \right.$$

5. Application to problems with non-local constraints

Let Ω be a bounded domain in \mathbf{R}^N , $1 \leq N < \infty$, with smooth boundary $\Gamma := \partial\Omega$, and let $X := W^{1,p}(\Omega)$ or $W_0^{1,p}(\Omega)$, $1 < p < \infty$. Let $a_i(x, \eta, \xi)$, $i = 1, 2, \dots, N$, and β be as in the previous section; conditions (4.1), (4.2) and (4.4) are satisfied as well. Furthermore let $k_c(\cdot)$ be a Lipschitz continuous function on \mathbf{R} with bounded Lipschitz continuous derivative $k'_c(\cdot)$ on \mathbf{R} ; condition (4.11) is satisfied as well.

Given a singlevalued compact mapping $\Lambda : X \rightarrow X$, we define constraint sets $K(v)$ in X by

$$K(v) := \{w \in X; w \geq k_c(\Lambda v) \text{ a.e. on } \Omega\}, \quad \forall v \in X. \quad (5.1)$$

Clearly, $K(v)$ is non-empty, closed and convex in X for every $v \in X$.

Lemma 5.1. *The family $\{K(v); v \in X\}$ given by (5.1) and the set $G_0 := \{k^*\}$ satisfy conditions (K'1) and (K2).*

Proof. Assume that $v_n \rightarrow v$ weakly in X and let w be any function in $K(v)$, namely $w \geq k_c(\Lambda v)$ a.e. on Ω . We note that $\Lambda v_n \rightarrow \Lambda v$ in X and hence $k_c(\Lambda v_n) \rightarrow k_c(\Lambda v)$ in X . Putting $w_n = w - k_c(\Lambda v) + k_c(\Lambda v_n)$, we see that $w_n \in K(v_n)$, $w_n \rightarrow w$ in X . Thus (K'1) is verified. Next, assume that $w_n \in K(v_n)$, $w_n \rightarrow w$ weakly in X and $v_n \rightarrow v$ weakly in X . Then $w_n \rightarrow w$ and $v_n \rightarrow v$ in $L^p(\Omega)$ as well as $k_c(\Lambda v_n) \rightarrow k_c(\Lambda v)$ in $L^p(\Omega)$. Hence $w \geq k_c(\Lambda v)$ a.e. on Ω , that is $w \in K(v)$. Thus (K2) is obtained. \diamond

For the same mapping $\tilde{A} := \tilde{A}_0 + \beta$ as in the previous section, all the conditions of Theorem 2.2 are verified. Therefore, given a function $f \in L^q(\Omega)$, $\frac{1}{p} + \frac{1}{q} = 1$, there exists a function $u \in X$ and $u^* \in L^q(\Omega)$ such that

$$\left\{ \begin{array}{l} u \in X, u \geq k_c(\Lambda u) \text{ a.e. on } \Omega, u^* \in \beta(u) \text{ a.e. on } \Omega, \\ \sum_{i=1}^N \int_{\Omega} a_i(x, u, \nabla u)(u_{x_i} - v_{x_i}) dx + \int_{\Omega} u^*(u - w) dx \leq \int_{\Omega} f(u - w) dx, \\ \forall w \in X \text{ with } w \geq k_c(\Lambda u) \text{ a.e. on } \Omega. \end{array} \right. \quad (5.2)$$

Next, consider typical examples of the mapping Λ .

(Example 5.1) Consider the case when $p = 2$ and $X := W^{1,2}(\Omega)$. Let ν be a positive number. Then, for each $v \in X$, the boundary value problem

$$-\nu \Delta v + v = u \text{ in } \Omega, \quad \frac{\partial v}{\partial n} = 0 \text{ on } \Gamma, \quad (5.3)$$

has a unique solution v in $W^{2,2}(\Omega)$. Now, we define $\Lambda : X \rightarrow X$ by $v = \Lambda u$, where v is the solution of (5.3) for $u \in X$. Since Λ is bounded and linear from X into $W^{2,2}(\Omega)$, we see that Λ is compact from X into itself; in fact, $\Lambda = (I - \nu \Delta)^{-1}$. In this case, given $f \in L^2(\Omega)$, the quasi-variational inequality (5.2) with $p = 2$ is a system to find $u \in W^{1,2}(\Omega)$ with $u^* \in L^2(\Omega)$ and $v \in W^{2,2}(\Omega)$ such that

$$\left\{ \begin{array}{l} u \geq k_c(v) \text{ a.e. on } \Omega, u^* \in \beta(u) \text{ a.e. on } \Omega, \\ \sum_{i=1}^N \int_{\Omega} a_i(x, u, \nabla u)(u_{x_i} - w_{x_i}) dx + \int_{\Omega} u^*(u - w) dx \leq \int_{\Omega} f(u - w) dx, \\ \forall w \in X \text{ with } w \geq k_c(v) \text{ a.e. on } \Omega, \\ -\nu \Delta v + v = u \text{ a.e. on } \Omega, \\ \frac{\partial v}{\partial n} = 0 \text{ a.e. on } \Gamma, \end{array} \right. \quad (5.4)$$

and by virtue of Theorem 2.2 the above system has at least one solution $\{u_\nu, v_\nu\}$ for each $\nu \in (0, 1]$. Also, see [12] for a related non-local quasi-variational inequality.

Proposition 5.1. *Assume that the space dimension is one, $p = 2$ and $\Omega = (0, 1)$. Let $\{u_\nu, v_\nu\}_{\nu \in (0, 1]}$ be a family of solutions of (5.4). Then $\{u_\nu, v_\nu\}$ is bounded in $W^{1,2}(\Omega) \times W^{1,2}(\Omega)$, and for any weak limit u of $\{u_\nu\}$ as $\nu \downarrow 0$ is a solution of the interior obstacle problem (4.14).*

Proof. By testing k^* of (4.11) in quasi-variational inequality in (5.4) with $u = u_\nu$, it is easy to see that $\{u_\nu\}_{\nu \in (0, 1]}$ is bounded in $W^{1,2}(0, 1)$, hence is relatively compact in $C([0, 1])$. Now, let u be any weak limit of $\{u_\nu\}$ and choose a sequence ν_n with $\nu_n \downarrow 0$ so that $u_n := u_{\nu_n} \rightarrow u$ weakly in $W^{1,2}(0, 1)$, hence $u_n := u_{\nu_n} \rightarrow u$ uniformly on $[0, 1]$, as $n \rightarrow \infty$. Putting $v_n := v_{\nu_n}$ and multiplying $-\nu_n \Delta v_n + v_n = u_n$ by v_n and $-\Delta v_n$, we obtain

$$\begin{aligned} \int_0^1 |\nabla v_n|^2 dx + \int_0^1 |v_n|^2 dx &\leq \int_0^1 |v_n| |u_n| dx, \\ \nu_n \int_0^1 |\Delta v_n|^2 dx + \int_0^1 |\nabla v_n|^2 dx &\leq \int_0^1 |\nabla v_n| |\nabla u_n| dx, \end{aligned}$$

whence $\{\nu_n |\Delta v_n|_{L^2(0,1)}^2\}$ is bounded and $\{v_n\}$ is bounded in $W^{1,2}(0, 1)$. Since $v_n - u_n = \nu_n \Delta v_n \rightarrow 0$ in $L^2(0, 1)$ and $u_n \geq k_c(v_n)$, it follows from the above estimates that

$$v_n \rightarrow u \text{ weakly in } W^{1,2}(\Omega) \text{ and uniformly on } [0, 1] \quad (5.5)$$

as well as

$$k_c(v_n) \rightarrow k_c(u) \text{ uniformly on } [0, 1], \quad u \geq k_c(u) \text{ on } (0, 1). \quad (5.6)$$

Let w be any function with $w \geq k_c(u)$ on $(0, 1)$. Given $\varepsilon > 0$, choose a positive integer n_ε so that

$$w + \varepsilon \geq k_c(u) + \varepsilon \geq k_c(v_n), \quad u + \varepsilon \geq k_c(u) + \varepsilon \geq k_c(v_n), \quad \text{on } (0, 1), \quad \forall n \geq n_\varepsilon; \quad (5.7)$$

this is possible on account of (5.5) and (5.6). Now, take $u + \varepsilon$ as w the quasi-variational inequality in (5.4) which u_n satisfies, to get

$$\int_0^1 a_1(x, u_n, u_{n,x})(u_{n,x} - u_x) dx + \int_0^1 u_n^*(u_n - u - \varepsilon) dx \leq \int_0^1 f(u_n - u - \varepsilon) dx,$$

where $u_n^* \in L^2(0, 1)$ is a function satisfying $u_n^* \beta(u_n)$ a.e. on $(0, 1)$. Then we have

$$H := \limsup_{n \rightarrow \infty} \int_0^1 a_1(x, u_n, u_{n,x})(u_{n,x} - u_x) dx + \int_0^1 u_n^*(u_n - u) dx \leq M_0 \varepsilon,$$

where M_0 is a positive constant. Since $\varepsilon > 0$ is arbitrary, it holds that $H \leq 0$. This inequality implies by the pseudo-monotonicity property that

$$\lim_{n \rightarrow \infty} \int_0^1 a_1(x, u_n, u_{n,x})(u_{n,x} - u_x) dx + \int_0^1 u_n^*(u_n - u) dx = 0 \quad (5.8)$$

and

$$\lim_{n \rightarrow \infty} \int_0^1 a_1(x, u_n, u_{n,x}) \tilde{w}_x dx + \int_0^1 u_n^* \tilde{w} dx = \int_0^1 a_1(x, u, u_x) \tilde{w}_x dx + \int_0^1 u^* \tilde{w} dx, \quad (5.9)$$

$$\forall \tilde{w} \in W^{1,2}(0,1),$$

where $u^* \in L^2(0,1)$ with $u^* \in \beta(u)$ a.e. on $(0,1)$. Going back to the quasi-variational inequality (5.4) with $\nu = \nu_n$ and $u = u_n$ and $v = v_n$ for any $n \geq n_\varepsilon$, we obtain by (5.7)

$$\int_0^1 a_i(x, u_n, u_{n,x})(u_{n,x} - w_x)dx + \int_0^1 u_n^*(u_n - w - \varepsilon)dx \leq \int_0^1 f(u_n - w - \varepsilon)dx$$

for all $w \in W^{1,2}(0,1)$ with $w \geq k_c(u)$ on $(0,1)$. Letting $n \rightarrow \infty$ in this equality and using (5.8) and (5.9), we see that

$$H(u, w) := \int_0^1 a_i(x, u, u_x)(u_x - w_x)dx + \int_0^1 u^*(u - w)dx - \int_0^1 f(u - w)dx \leq M_0\varepsilon$$

for all $w \in W^{1,2}(0,1)$ with $w \geq k_c(u)$ on $(0,1)$. By the arbitrariness of $\varepsilon > 0$ we conclude that $H(u, w) \leq 0$ and u is a solution of quasi-variational inequality (4.14). \diamond

(Example 5.2) The second example of Λ is given as an integral operator as follows. Let $\rho(\cdot, \cdot, \cdot)$ be a smooth function on $\mathbf{R}^N \times \mathbf{R}^N \times \mathbf{R}$ such that

$$|\rho(x, y, r)| \leq c_4|r| + c'_4, \quad \forall x, y \in \mathbf{R}^N, \quad \forall r \in \mathbf{R},$$

where c_4 and c'_4 are positive constants. Then we define $\Lambda : X := W^{1,p}(\Omega) \rightarrow X$ by

$$(\Lambda v)(x) = \int_\Omega \rho(x, y, v(y))dy, \quad \forall x \in \Omega, \quad \forall v \in X.$$

Clearly Λ is compact from X into itself.

Now we give some concrete examples of integral operator Λ .

(Case 1) We consider as Λ the usual convolution operator by means of mollifier ρ_ε with real parameter $0 < \varepsilon < 1$. Let

$$\rho_\varepsilon(x, y) := \frac{1}{\varepsilon^N} \rho_0\left(\frac{x-y}{\varepsilon}\right), \quad \forall x, y \in \mathbf{R}^N,$$

where

$$\rho_0(x) := \begin{cases} c \cdot \exp\left\{-\frac{1}{1-|x|^2}\right\}, & \text{if } |x| < 1, \\ 0, & \text{otherwise,} \end{cases}$$

the constant $c > 0$ being chosen so that $\int_{\mathbf{R}^N} \rho_0 dx = 1$. Then, the mapping $\Lambda = \Lambda_\varepsilon$ is given by

$$(\Lambda_\varepsilon v)(x) = \int_\Omega \rho_\varepsilon(x-y)v(y)dy, \quad \forall v \in X, \quad (5.10)$$

as well as the constraint set $K(v) = K_\varepsilon(v)$ is of the form:

$$K_\varepsilon(v) = \left\{ w \in X; w \geq k_c\left(\int_\Omega \rho_\varepsilon(\cdot - y)v(y)dy\right) \text{ a.e. on } \Omega \right\}.$$

According to the above existence result, for each $\varepsilon > 0$ problem (5.2) with $\Lambda := \Lambda_\varepsilon$ has at least one solution u_ε ($\in K_\varepsilon(u_\varepsilon)$). For the family $\{u_\varepsilon\}$ of solutions we see by taking $w = k^*$

in inequality (5.2) that $\{u_\varepsilon\}_{0 < \varepsilon < 1}$ is bounded in X . It is a quite interesting question what happens as $\varepsilon \rightarrow 0$.

Proposition 5.2. *Assume that the space dimension is one and $\Omega = (0, 1)$. Let $\{u_\varepsilon\}_{\varepsilon \in (0, 1]}$ be a family of solutions of (5.2) with $\Lambda = \Lambda_\varepsilon$. Then $\{u_\varepsilon\}$ is bounded in $W^{1,p}(0, 1)$ and any weak limit u of $\{u_\varepsilon\}$ as $\varepsilon \downarrow 0$ is a solution of the interior obstacle problem (4.14).*

Proof. We take $w \equiv k^*$ in (5.2) with $u = u_\varepsilon$ and $\Lambda = \Lambda_\varepsilon$ to get

$$\int_0^1 a_1(x, u_\varepsilon, u_{\varepsilon,x}) u_{\varepsilon,x} dx + \int_0^1 u_\varepsilon^*(u_\varepsilon - k^*) dx \leq \int_0^1 f(u_\varepsilon - k^*) dx.$$

It is easy to see from this inequality that $\{u_\varepsilon\}$ is bounded in $W^{1,p}(0, 1)$. Now let u be any weak limit of $\{u_\varepsilon\}$ in $W^{1,p}(0, 1)$ and choose a sequence $\varepsilon_n \downarrow 0$ so that $u_n := u_{\varepsilon_n} \rightarrow u$ weakly in $W^{1,p}(0, 1)$ and uniformly on $[0, 1]$. In this case we have that $\Lambda_{\varepsilon_n} u_n \rightarrow u$ uniformly on $[0, 1]$ and $u \geq k_c(u)$ on $(0, 1)$. Therefore, in a way similar to that of Proposition 5.1, we can prove that u is a solution of the quasi-variational inequality (4.14). \diamond

(Case 2) Consider $\Omega = (0, 1)$ and define $\rho(x, y)$ by

$$\rho(x, y) := \begin{cases} s(x - y), & \text{if } y \in [x, 1), \\ 0, & \text{if } y \in (0, x), \end{cases} \quad (5.11)$$

where $s(\cdot)$ is a non-negative smooth function on \mathbf{R} such that $s(x) = 0$ for $x \leq 0$. Then, we define $\Lambda : X := W^{1,p}(0, 1) \rightarrow X$ by

$$(\Lambda v)(x) = \int_0^1 s(x - y) v(y) dy, \quad \forall v \in X.$$

where $k_c(\cdot)$ is as before. In this case, by the definition of quasi-variational inequality any solution u of (5.2) has to satisfy that

$$u(x) \geq k_c \left(\int_0^1 s(x - y) u(y) dy \right), \quad \text{a.e. } x \in (0, 1). \quad (5.12)$$

As is easily understood, the behaviour of u satisfying (5.12) is controlled by that of $s(\cdot)$, and some suitable classes of functions $s(\cdot)$ are expected to play a role of an effective control space, for instance, in optimal design problems. In this sense, it is interesting to derive necessary conditions of u in order to satisfy (5.12). For instance, it is easily checked from (5.12) that $u(1) \geq k_c(0)$, because, by (5.11), $s(1 - y) = 0$ for all $y \in (0, 1)$. This means that it may be possible to control the solution u of problem (5.2) by appropriate choice of convolution kernel $s(\cdot)$.

(Case 3) In case $\rho(x, y)$ is constant $|\Omega|$ on $\mathbf{R}^N \times \mathbf{R}^N$, where $|\Omega|$ is the volume of Ω , we define

$$(\Lambda v)(x) = \frac{1}{|\Omega|} \int_\Omega |v(y)| dy, \quad \forall x \in \Omega, \quad \forall v \in X := W^{1,p}(\Omega).$$

In this case, we see that Λ is a continuous mapping from the weak topology of X into \mathbf{R} , which is regarded as a compact mapping from X into itself.

References

1. C. Baiocchi and A. Capelo, Variational and Quasivariational Inequalities, John Wiley and Sons, Chichester-New York-Brisbane-Toronto-Singapore, 1984.
2. A. Bensoussan and J.L. Lions, Nouvelle formulation de problèmes de contrôle impulsionnel et applications, C. R. Acad. Sci. Paris Sér. A, 276(1973), 1189-1192.
3. A. Bensoussan, M. Goursat, J.L. Lions, Contrôle impulsionnel et inéquations quasi-variationnelles stationnaires, C. R. Acad. Sci. Paris Ser. A, vol.276(1973), 1279-1284.
4. H. Brézis, Équations et inéaires dans les espaces vectoriels en dualité, Ann. Inst. Fourier, Grenoble, 18 (1968), 115-175.
5. F.E. Browder, Nonlinear monotone operators and convex sets in Banach spaces, Bull. Amer. Math. Soc., 71(1965), 780-785.
6. J.L. Joly and U. Mosco, Sur les inéquations quasi-variationnelles, C. R. Acad. Sci. Paris Ser. A, 279(1974), 499-502.
7. J.L. Joly and U. Mosco, A propos de l'existence et de la régularité des solutions de certaines inéquations quasi-variationnelles, J. Functional Analysis, 34(1979), 107-137.
8. N. Kenmochi, Nonlinear operators of monotone type in reflexive Banach spaces and nonlinear perturbations, Hiroshima Math. J., 4(1974), 229-263.
9. T. Laetsch, A uniqueness theorem for elliptic quasi-variational inequalities, J. Functional Analysis, 18(1975), 286-287.
10. J.L. Lions and G. Stampacchia, Variational inequalities, Comm. Pure Appl. Math., 20(1967), 493-519.
11. U. Mosco, Convergence of convex sets and of variational inequalities, Advances Math., 3(1969), 510-585.
12. N. Nassif and K. Malla, Étude de l'existence de la solution d'une inégalité quasi-variationnelle apparaissant dans la théorie des semi-conducteurs, C. R. Acad. Sci. Paris Serie 1, 294(1982), 119-122.
13. L. Tartar, Inéquations quasi-variationnelles abstraites, C. R. Acad. Sci. Paris Ser. A, 278(1974), 1193-1196.