

# Jacobi type formula for Matsumoto hyperelliptic curves

Hironori Shiga (Graduate School of Mathematics and Physics, Chiba Univ.)

On the occasion of the final lecture at Chiba university

## 1 Introduction

This is a continuation of our previous work “ A variant of Jacobi type formula for Picard curves ”([M-S]). Here we start from the algebraic curve  $w^4 = z^2(z-1)^2(z-\lambda_1)(z-\lambda_2)$  with two complex parameters  $\lambda_1, \lambda_2$ . Generically it is a hyperelliptic curve of genus three. This curve has been studied by K. Matsumoto [M] ( and see also [K-S] ) in detail. We call it Matsumoto hyperelliptic curve. In this article we show a Jacobi type formula of two variables that connects the theta function and the hypergeometric function via the modular function for the Matsumoto hyperelliptic curves defined on the two dimensional hyperball. Our main theorem is stated in (2.6). As an application, we give a new proof for the extended Gauss arithmetic-geometric mean theorem discovered in [K-S]. We consider this is one aspect of various natures of the  $K3$  modular function that was initiated by the author in 1977 (see [S1], [S2]). And he believes that this result will be understood in a more general context in the future.

## 2 Statement of Main Theorem

First recall the classical Jacobi formula. Set

$$\lambda(\tau) = \frac{\vartheta_{01}^4(\tau)}{\vartheta_{00}^4(\tau)} \quad (2.1)$$

for  $\tau \in \mathbf{H} = \{\text{Im } \tau > 0\}$ , where  $\vartheta_{jk}$  indicates the Jacobi theta constant

$$\vartheta_{jk}(\tau) = \sum_{n \in \mathbb{Z}} \exp[\pi i(n + \frac{j}{2})^2 \tau + 2\pi i(n + \frac{j}{2})\frac{k}{2}].$$

**Theorem 2.1.** (Classical Jacobi Formula, see [J] p.235) *Under the relation (2.1) we have*

$$\vartheta_{00}^2(\tau) = F\left(\frac{1}{2}, \frac{1}{2}, 1; 1-\lambda\right) = \frac{1}{\pi} \int_1^\infty \frac{dz}{\sqrt{z(z-1)(z-(1-\lambda))}}. \quad (2.2)$$

We have a two dimensional counterpart of this theorem.

Set

$$\mathbf{B} = \{(u, v) \in \mathbf{C}^2 : 2 \text{Im } v - |u|^2 > 0\}, \quad (2.3)$$

and set

$$\Omega(u, v) = \begin{pmatrix} v + iu^2/2 & -u^2/2 & -iu \\ -u^2/2 & v - iu^2/2 & u \\ -iu & u & i \end{pmatrix}.$$

Recall the Riemann theta constant

$$\vartheta \begin{bmatrix} a \\ b \end{bmatrix} (\Omega) = \sum_{n \in \mathbb{Z}^3} \exp[\pi i(n+a)\Omega^t(n+a) + 2\pi i(n+a)^t b],$$

here  $a$  and  $b$  are row vectors in  $\mathbf{Q}^3$ , and  $\Omega$  is a variable on the Siegel upper half space of degree 3. Let us denote

$$\vartheta \begin{bmatrix} a \\ b \end{bmatrix} (\Omega(u, v)) = \Theta \begin{bmatrix} a \\ b \end{bmatrix} (u, v) \text{ for } (u, v) \in \mathbf{B}.$$

Set

$$(\lambda_1, \lambda_2) = \left( \frac{\Theta^4 \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \end{bmatrix} (u, v)}{\Theta^4 \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} (u, v)}, \frac{\Theta^4 \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix} (u, v)}{\Theta^4 \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} (u, v)} \right). \quad (2.4)$$

**Theorem 2.2.** (Main Theorem: Extended Jacobi formula)

For the Appell hypergeometric function

$$F_1\left(\frac{1}{2}, \frac{1}{4}, \frac{1}{4}, 1; x, y\right) = \sum_{m, n \geq 0} \frac{(\frac{1}{2}, m+n) (\frac{1}{4}, m) (\frac{1}{4}, n)}{(1, m+n) m! n!} x^m y^n, \quad (2.5)$$

where  $(a, n) = a(a+1) \cdots (a+n-1)$  for  $n \geq 1$  and  $(a, 0) = 1$ , we have

$$\Theta \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} (u, v) = \vartheta_{00}(i) F_1\left(\frac{1}{2}, \frac{1}{4}, \frac{1}{4}, 1; 1 - \lambda_1, 1 - \lambda_2\right), \quad (2.6)$$

with  $(\lambda_1, \lambda_2)$  given by (2.4).

**Remark 2.1.** Here we set  $(\lambda_1, \lambda_2)$  in the real triangle  $\{(\lambda_1, \lambda_2) \in (0, 1) \times (0, 1) : \lambda_1 < \lambda_2\}$ . Then we can find the variables  $(u, v) \in \mathbf{B}$  with the relation (2.4). At first we have an exact equality only for these variables, next we can make the analytic continuations of both sides on the whole domain  $\mathbf{B}$ .

**Remark 2.2.** If we put  $u = 0$ , then by using (2.4) we have  $\lambda_1 = \lambda_2$ . In this case, our main theorem coincides with the classical Jacobi formula (2.2).

## 3 Geometric back ground

### 3.1 Matsumoto hyperelliptic modular function

Set

$$C(\lambda) = C(\lambda_1, \lambda_2) : w^4 = z^2(z-1)^2(z-\lambda_1)(z-\lambda_2) \quad (3.1)$$

for  $\lambda = (\lambda_1, \lambda_2) \in \Lambda = \{(\lambda_1, \lambda_2) \in \mathbf{C}^2 : \lambda_1 \lambda_2 (\lambda_1 - 1)(\lambda_2 - 1)(\lambda_1 - \lambda_2) \neq 0\}$ . It is a hyperelliptic curve of genus 3. The differentials

$$dz/w, z(z-1)dz/w^3, z^2(z-1)dz/w^3$$

form a basis of holomorphic 1-forms on  $C(\lambda)$ .

For the moment suppose that  $\lambda_1$  and  $\lambda_2$  are real and satisfy  $0 \leq \lambda_1 < \lambda_2 < 1$ . Let  $\rho$  be the automorphism  $(z, w) \mapsto (z, \sqrt{-1}w)$  of  $C(\lambda)$ .

Set

$$a_1(\lambda_1, \lambda_2) = 2 \int_{-\infty}^0 \frac{dz}{w}, \quad a_2(\lambda_1, \lambda_2) = 2i \int_1^{+\infty} \frac{dz}{w}, \quad a_5(\lambda_1, \lambda_2) = (1-i) \int_{\lambda_1}^{\lambda_2} \frac{dz}{w},$$

here we used the real positive branch of  $w$  for  $a_1, a_2$  and  $-iw > 0$  for  $a_5$ . They are periods of  $C(\lambda)$ .

[Fact 1]  $a_1(\lambda_1, \lambda_2)$ ,  $a_2(\lambda_1, \lambda_2)$ ,  $a_5(\lambda_1, \lambda_2)$  satisfy the differential equation  $E_1(\frac{1}{2}, \frac{1}{4}, \frac{1}{4}, 1)$  of the Appell hypergeometric function in (2.5), and they form a basis of the space of solutions.

We define the Schwarz map  $\Phi : \Lambda \rightarrow \mathbf{C}^2$  for  $E_1(\frac{1}{2}, \frac{1}{4}, \frac{1}{4}, 1)$  by

$$\Phi : (\lambda_1, \lambda_2) \mapsto (u, v) = (a_5/a_1, a_2/a_1) \in \mathbf{C}^2.$$

[Fact 2] The image of  $\Phi$  is contained in the hyperball  $\mathbf{B}$  of (2.3), and it is open dense in  $\mathbf{B}$ . The map  $\Phi$  has a continuation on  $\mathbf{P}^1 \times \mathbf{P}^1 - \{(0, 0), (1, 1), (\infty, \infty)\}$  and the image is equal to  $\mathbf{B}$ .

[Fact 3] The fundamental group  $\pi_1(\Lambda, *)$  induces the monodromy group  $G$  of  $\Phi$  acting on  $\mathbf{B}$ . The following five transformations give a generator system of  $G$ :

$$\begin{aligned} g_1 &= \begin{pmatrix} 1 & 0 & 0 \\ 1+i & 1 & 1-i \\ -1-i & 0 & i \end{pmatrix}, \\ g_2 &= \begin{pmatrix} 2+i & -1-i & -1-i \\ 1+i & -i & -1-i \\ 1-i & -1+i & i \end{pmatrix}, \\ g_3 &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \\ g_4 &= \begin{pmatrix} i & 1-i & 1-i \\ 0 & i & 0 \\ 0 & -1-i & -1 \end{pmatrix}, \\ g_5 &= \begin{pmatrix} 2+i & -1-i & 1-i \\ 1+i & -i & 1-i \\ -1-i & 1+i & i \end{pmatrix}, \end{aligned}$$

where  $g_i$  ( $i = 1, 2, 3, 4, 5$ ) acts on  ${}^t(1, v, u)$  from left. Their orders are 4, 4, 2, 4, 4 and eigen values are

$$\{1, 1, i\}, \{1, 1, i\}, \{1, 1, -1\}, \{-1, i, i\}, \{1, 1, i\},$$

respectively (the original article contains a printing error for  $g_5$ ).

[Fact 4] The Schwarz map  $\Phi$  induces a biholomorphic correspondence

$$\mathbf{P}^1 \times \mathbf{P}^1 \cong \overline{\mathbf{B}/G},$$

and three points  $(0, 0), (1, 1), (\infty, \infty)$  correspond to the boundary  $\overline{\mathbf{B}/G} - \mathbf{B}/G$ . We have the representatives

$$\Phi(0, 0) = (0, 0), \Phi(1, 1) = (0, -1), \Phi(\infty, \infty) = (0, i\infty)$$

on  $\partial\mathbf{B}$ .

[Fact 5] We can realize  $\mathbf{B}$  in the form:

$$\begin{aligned} \{\xi = [\xi_0, \xi_1, \xi_2] \in \mathbf{P}^2 : \xi H^t \bar{\xi} < 0\}, \\ H = \begin{pmatrix} 0 & -i & 0 \\ i & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}. \end{aligned}$$

The map

$$\Omega(u, v) = \begin{pmatrix} v + iu^2/2 & -u^2/2 & -iu \\ -u^2/2 & v - iu^2/2 & u \\ -iu & u & i \end{pmatrix}$$

gives an modular embedding of  $\mathbf{B}$  into the Siegel upper half space of degree 3. Set

$$G_0 = U(2, 1, \mathbf{Z}[i]) = \{g \in GL(3, \mathbf{Z}[i]) : gH^t \bar{g} = H\}.$$

$G_0$  becomes to be a restriction of  $Sp(6, \mathbf{Z})$  on  $\Omega(\mathbf{B})$ .

[Fact 6] Set

$$g_6 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & i \end{pmatrix}.$$

And let  $G$  be the monodromy group of  $E_1(\frac{1}{2}, \frac{1}{4}, \frac{1}{4})$ . Then the group generated by  $G$  and  $g_6$  together with  $i = i \cdot \text{id}$  becomes to be the congruence subgroup

$$G(1+i) := \{g \in G_0 : g \equiv \text{id} \pmod{(1+i)}\}.$$

[Fact 7] We set the following theta constants:

$$\begin{aligned} \Theta_{xN} &= \Theta \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{bmatrix} (u, v), & \Theta_{xD} &= \Theta \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{bmatrix} (u, v), \\ \Theta_{yN} &= \Theta \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix} (u, v), & \Theta_{yD} &= \Theta \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{bmatrix} (u, v), \\ \Theta_{h1} &= \Theta \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} (u, v), & \Theta_Z &= \Theta \begin{bmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix} (u, v), & \Theta_{h3} &= \Theta \begin{bmatrix} 0 & 0 & 0 \\ 1 & 1 & 0 \end{bmatrix} (u, v), \end{aligned} \quad (3.2)$$

here we used the same notation for the Riemann theta constant defined in the previous section. We have the following algebraic relations (see [K-S]):

$$\begin{aligned} \Theta_{h3}(u, v) &= \Theta_{h1}(u, v) \frac{\Theta_{xD}(u, v)\Theta_{yD}(u, v)}{\Theta_{xN}(u, v)\Theta_{yN}(u, v)} \\ \Theta_Z(u, v) &= \Theta_{h1}(u, v) \frac{1}{\Theta_{xN}(u, v)\Theta_{yN}(u, v)} \sqrt{\frac{1}{2} \left( \Theta_{xD}^2(u, v)\Theta_{yN}^2(u, v) + \Theta_{yD}^2(u, v)\Theta_{xN}^2(u, v) \right)}. \end{aligned}$$

[Fact 8] ::Matsumoto hyperelliptic theta formula (see [M] and [K-S]):

It holds

$$(\lambda_1, \lambda_2) = \left( \frac{\Theta^4 \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \end{bmatrix} (u, v)}{\Theta^4 \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} (u, v)}, \frac{\Theta^4 \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix} (u, v)}{\Theta^4 \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} (u, v)} \right). \quad (3.3)$$

### 3.2 modular behavior of the theta constants

**Proposition 3.1.** *Every fourth power of the theta constants listed in (3.2), except  $\Theta_Z$ , is a modular form of weight 4 with respect to  $G$ , namely it satisfies*

$$\Theta^4 \begin{bmatrix} a \\ b \end{bmatrix} (g \circ (u, v)) = (p_1 + p_2v + p_3u)^4 \Theta^4 \begin{bmatrix} a \\ b \end{bmatrix} (u, v), \quad g = \begin{pmatrix} p_1 & p_2 & p_3 \\ \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \end{pmatrix} \in G.$$

**Proposition 3.2.** (see Prop. 2.4 in [K-S])

*The vector space of modular forms of weight 4 with respect to  $G$  is generated by  $\Theta_{xN}^4, \Theta_{yN}^4, \Theta_{h1}^4, \Theta_{h3}^4$ .*

**Remark 3.1.** *More precisely we have*

$$\begin{aligned} \Theta_{h1}(g_2(u, v)) &= i(2+i+(-1-i)v+(-1-i)u)\Theta_{h1}(u, v), \\ \Theta_{h1}(g_5(u, v)) &= i(2+i+(-1-i)v+(1-i)u)\Theta_{h1}(u, v). \end{aligned}$$

*We shall use these formula in our proof of Main Theorem.*

## 4 Proof of Main Theorem

**Lemma 4.1.** *We have*

$$F_1\left(\frac{1}{2}, \frac{1}{4}, \frac{1}{4}, 1; 1 - \lambda_1, 1 - \lambda_2\right) = \frac{1}{\pi} \int_{-\infty}^0 \frac{dz}{w} = \frac{1}{2\pi} a_1(\lambda_1, \lambda_2) \quad (0 < \lambda_1, \lambda_2 < 1). \quad (4.1)$$

Proof]. According to [App] we have

$$F_1\left(\frac{1}{2}, \frac{1}{4}, \frac{1}{4}, 1; s, t\right) = \frac{1}{\pi} \int_1^{\infty} z'^{-1/2} (z' - 1)^{-1/2} (z' - s)^{-1/4} (z' - t)^{-1/4} dz' \quad (|s|, |t| < 1).$$

By changing the variable  $z' = 1 - z$  we obtain the required equality.

q.e.d.

**Lemma 4.2.**  $\Theta^4 \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} (u, v)$  and  $F_1(\frac{1}{2}, \frac{1}{4}, \frac{1}{4}, 1; 1 - \lambda_1, 1 - \lambda_2)^4$  are modular forms of weight 4 with respective to  $G$ .

Proof]. The first statement is a direct consequence of Prop. 3.1. Using Lemma 4.1 the second statement is derived from the monodromy action

$$\begin{pmatrix} a_1(g(u, v)) \\ a_2(g(u, v)) \\ a_5(g(u, v)) \end{pmatrix} = \begin{pmatrix} p_1 & p_2 & p_3 \\ \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \end{pmatrix} \begin{pmatrix} a_1(u, v) \\ a_2(u, v) \\ a_5(u, v) \end{pmatrix} \quad \text{with } g = \begin{pmatrix} p_1 & p_2 & p_3 \\ \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \end{pmatrix} \in G.$$

q.e.d.

**Lemma 4.3.** *The modular form  $a_1(u, v)$  vanishes along the divisor  $V_1$  on  $\mathbf{B}/G$  represented by  $\{u + v = 1\} + \{iu + v = 1\}$ .*

Proof of Main Theorem]. The differential  $\frac{dz}{w}$  on  $C(\lambda_1, \lambda_2)$  is a holomorphic differential for every point  $(\lambda_1, \lambda_2)$  on the affine part  $\mathbf{C} \times \mathbf{C}$  except two points  $(0, 0)$  and  $(1, 1)$ . So  $a_1(u, v)$  does not vanish on the affine part. And we can see that  $D_\lambda = \{\lambda_1 = \infty\} + \{\lambda_2 = \infty\}$  is the zero divisor of  $a_1(\lambda_1, \lambda_2)$ . By the direct calculation of the integrals  $a_1, a_2, a_5$  on  $D_\lambda$ , we know that  $\Phi(D_\lambda) = V_1$ . Hence the zero divisor of  $a_1(u, v)$  is equal to  $V_1$ .

According to Remark 3.1, we have

$$\Theta \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} (g_2(u, v)) = i(2 + i + (-1 - i)u + (-1 - i)v) \Theta \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} (u, v),$$

here

$$g_2(u, v) = \left( \frac{(1 - i) + iu - (1 - i)v}{(2 + i) - (1 + i)u - (1 + i)v}, \frac{(1 + i) - (1 + i)u - iv}{(2 + i) - (1 + i)u - (1 + i)v} \right).$$

And we have  $g_2(u, v) = (u, v)$  if and only if  $u = 1 - v$ . So it holds

$$\Theta \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} (u, v) = i \Theta \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} (u, v)$$

for  $(u, v) \in \mathbf{B}$  with  $u + v = 1$ . Namely we have  $\Theta \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} (u, v) = 0$  on the divisor  $\{u = 1 - v\} \subset \mathbf{B}$ .

By the same argument for  $g_5$ , we have  $\Theta \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} (u, v) = 0$  on the divisor  $\{u = -i(1 - v)\} \subset \mathbf{B}$ . Hence the zero divisor of  $\Theta_{h_1}(u, v)$  is greater than or equal to  $V_1$ .

Because  $a_1(u, v)^4$  and  $\Theta_{h_1}^4$  are modular forms with the same automorphic behavior, their zero divisors are linearly equivalent. So the zero divisor of  $\Theta_{h_1}(u, v)$  is equal to  $V_1$ . Hence we have

$$\Theta_{h_1}(u, v) = c \cdot a_1(u, v)$$

with some constant  $c$ . We can determine it by the calculation on  $u = 0$ . Consequently, we obtain the required equality.

q.e.d.

## 5 Application to the Extended Gauss AGM

It holds the following CM-isogeny formula (see [K-S]):

**Theorem 5.1.**

$$\begin{cases} \Theta_{h_1}((1+i)u, 2v) = \frac{1}{2}(\Theta_{h_1}(u, v) + \Theta_{h_3}(u, v)), \\ \Theta_{h_3}((1+i)u, 2v) = \Theta_Z(u, v), \\ \Theta_Z((1+i)u, 2v) = \sqrt[4]{\frac{1}{4}(\Theta_Z^2 + \Theta_{h_1}\Theta_{h_3})(\Theta_Z^2 + \frac{\Theta_{h_1}^2 + \Theta_{h_3}^2}{2})}. \end{cases} \quad (5.1)$$

Let  $(a, b, c)$  be a triple of positive real numbers. Set an AGM system:

$$(a', b', c') = \psi(a, b, c) = \left( \frac{1}{2}(a+b), c, \sqrt[4]{\frac{1}{4}(c^2+ab)(c^2+(a^2+b^2)/2)} \right). \quad (5.2)$$

We can define the common limit  $\lim_{n \rightarrow \infty} \psi^n(a, b, c)$ . And we denote it by  $M_{ext}(a, b, c)$ .

**Corollary 5.1.** (Extended Gauss AGM formula [K-S]) *Assume  $0 < x, y < 1$  and set  $\lambda_1 = 1 - x^2$ ,  $\lambda_2 = 1 - y^2$ . Then we have :*

$$\frac{1}{M_{ext}(1, \sqrt{xy}, \sqrt{(x+y)/2})} = \frac{1}{\pi} \int_1^\infty \frac{du}{\sqrt{u^2(u-1)^2(u-\lambda_1)(u-\lambda_2)}} = F_1\left(\frac{1}{2}, \frac{1}{4}, \frac{1}{4}, 1; \lambda_1, \lambda_2\right). \quad (5.3)$$

[Proof]. Put

$$x = \frac{\Theta_{xD}^2(u, v)}{\Theta_{xN}^2(u, v)}, y = \frac{\Theta_{yD}^2(u, v)}{\Theta_{yN}^2(u, v)}.$$

According to Fact 7, we have

$$(a, b, c) = \Theta_{h_1}(u, v)(1, \sqrt{xy}, \sqrt{\frac{1}{2}(x+y)}) = (\Theta_{h_1}(u, v), \Theta_{h_3}(u, v), \Theta_Z(u, v))$$

with

$$\begin{cases} b = \Theta_{h_1}(u, v) \frac{\Theta_{xD}(u, v)\Theta_{yD}(u, v)}{\Theta_{xN}(u, v)\Theta_{yN}(u, v)}, \\ c = \Theta_{h_1}(u, v) \frac{1}{\Theta_{xN}(u, v)\Theta_{yN}(u, v)} \sqrt{\frac{1}{2}(\Theta_{xD}^2(u, v)\Theta_{yN}^2(u, v) + \Theta_{yD}^2(u, v)\Theta_{xN}^2(u, v))}. \end{cases}$$

By using the isogeny formula (5.1) we have

$$\psi(a, b, c) = (\Theta_{h_1}((1+i)u, 2v), \Theta_{h_3}((1+i)u, 2v), \Theta_Z((1+i)u, 2v)).$$

Consequently we have

$$\Theta_{h_1}(u, v)M_{ext}(1, \sqrt{xy}, \sqrt{(x+y)/2}) = \lim_{n \rightarrow \infty} \psi^n(a, b, c) = \vartheta_{00}(i)(1, 1, 1).$$

According to our Main Theorem, we have

$$\vartheta_{00}(i) = \frac{\Theta_{h_1}(u, v)}{F_1\left(\frac{1}{2}, \frac{1}{4}, \frac{1}{4}, 1; 1-x^2, 1-y^2\right)}.$$

By conjunction of these two equalities we obtain the required equality (5.3).

q.e.d.

## References

- [App] P. Appell, *Sur les Fonctions hypergéométriques de plusieurs variables les polynomes d'Hermite et autres fonctions sphériques dans l'hyperspace*, Gauthier-Villars, Paris, 1925.
- [J] C.G.J. Jacobi, *Gesammelte Werke I*, Chelsea, New York, 1969.
- [K-S] K. Koike and H. Shiga, *An extended Gauss AGM and corresponding Picard modular forms*, J. Number Theory, 128(2008), 2029–2126.
- [M-S] K. Matsumoto and H. Shiga, *A variant of Jacobi type formula for Picard curves*, Tech. Report Chiba Univ., 2008.
- [M] K. Matsumoto, *On modular functions in 2 variables attached to a family of hyperelliptic curves of genus 3*, Annali S. N. S. di Pisa - Classe di Scienze, Ser. IV, vol. XVI, no. 4 (1989), p. 557–578.
- [S1] H. Shiga, *One attempt to the K3 modular function*, Several Complex Variables, Proc. Int. Conferences Cortona 1976-1977, S. N. S. Pisa (1978), 277–281.
- [S2] H. Shiga, *One attempt to the K3 modular function I*, Annali S. N. S. di Pisa - Classe di Scienze, Ser. IV, vol. VI, no. 4 (1979), p. 609 – 635.

Hironori Shiga,  
Graduate School of Mathematics and Physics, Chiba University,  
Inage-ku, Yayoi-cho 1-33, Chiba, 263-8522 Japan  
shiga@math.s.chiba-u.ac.jp