

# THE ORDER OF CONFORMAL AUTOMORPHISMS OF RIEMANN SURFACES OF INFINITE TYPE

EGE FUJIKAWA

ABSTRACT. Let  $R$  be a Riemann surface of infinite type such that the injectivity radius at any point in  $R$  is less than a positive constant  $M$ , and  $f$  a conformal automorphism of  $R$  fixing a compact subset in  $R$ . We show that the order of  $f$  is less than a certain constant depending on  $M$ .

## 1. INTRODUCTION

On a compact Riemann surface  $R$  of genus  $g \geq 2$ , it is known that the order of a conformal automorphism of  $R$  is not greater than  $2(2g + 1)$  (Wiman cf. [4, p.96]). Since the hyperbolic area of  $R$  is  $4\pi(g - 1)$ , the injectivity radius at any point in  $R$  is not greater than a constant depending only on  $g$ . This means that the order of a conformal automorphism of  $R$  is estimated by the supremum of the injectivity radii which is taken over all points in  $R$ . We extend this result to the case of Riemann surfaces which are not necessarily of finite type. That is, for any hyperbolic Riemann surface  $R$  such that the injectivity radius at any point in  $R$  is less than a positive constant  $M$ , if a conformal automorphism  $f$  of  $R$  fixes a compact subset in  $R$ , then the order of  $f$  is estimated by  $M$ . Note that, in the case that  $R$  has the non-abelian fundamental group, a conformal automorphism  $f$  of  $R$  fixes a compact subset on  $R$  if and only if  $f$  has the finite order.

## 2. MAIN THEOREMS

Let  $\mathbb{H}$  be the upper-half plane equipped with the hyperbolic metric  $d\lambda = |dz|/\text{Im}z$ . We say that a Riemann surface  $R$  is *hyperbolic* if it is represented by  $\mathbb{H}/\Gamma$  for a torsion-free Fuchsian group  $\Gamma$  acting on  $\mathbb{H}$ . The hyperbolic distance on  $\mathbb{H}$  or on  $R$  is denoted by  $d(\cdot, \cdot)$ . The *injectivity radius* at  $p \in R$  is the supremum of radii of embedded hyperbolic discs centered at  $p$ .

Before we state the main theorems, we note the following fact.

**Proposition 1.** *Let  $R = \mathbb{H}/\Gamma$ , where  $\Gamma$  is a Fuchsian group which is not necessarily torsion-free, and  $f$  a conformal automorphism of  $R$  with finite order  $n > 1$ . Then  $f$  fixes either a simple closed geodesic, a puncture, a point or a cone point on  $R$ .*

*Proof.* Let  $\tilde{f}$  be a lift of  $f$  to  $\mathbb{H}$  which is an element of  $\mathrm{PSL}_2(\mathbb{R})$ . Since  $f$  has the finite order  $n$ , we see that  $\tilde{f}^n$  belongs to  $\Gamma$  and that  $\tilde{f}^m$  ( $1 \leq m < n$ ) does not belong to  $\Gamma$ . If  $\tilde{f}^n$  is parabolic, then  $\tilde{f}$  is parabolic. Hence  $f$  fixes a puncture on  $R$ . If  $\tilde{f}^n$  is the identity, then  $\tilde{f}$  is elliptic with the fixed point  $\tilde{p} \in \mathbb{H}$ . Hence  $f$  fixes the point on  $R$  which is the projection of  $\tilde{p}$ . If  $\tilde{f}^n$  is elliptic, then  $\tilde{f}$  is elliptic with the fixed point  $\tilde{p} \in \mathbb{H}$ . Hence  $f$  fixes the cone point on  $R$  which is the projection of  $\tilde{p}$ . Further, if  $\tilde{f}^n$  is hyperbolic, then  $\tilde{f}$  is hyperbolic. Hence  $f$  fixes a closed geodesic  $c_*$  on  $R$ . In this case, we prove that  $f$  fixes either a simple closed geodesic, a puncture, a point or a cone point on  $R$ . We consider the quotient  $\hat{R} = R/\langle f \rangle$  by a cyclic group  $\langle f \rangle$  and its Fuchsian model  $\hat{\Gamma} = \langle \Gamma, \tilde{f} \rangle$ . Then  $\hat{c}_* = c_*/\langle f \rangle$  is a closed geodesic on  $\hat{R}$ . There exists a subset  $\hat{c}'$  of  $\hat{c}_*$  such that  $\hat{c}'$  is a non-trivial simple closed curve and it corresponds to a conjugacy class of an element  $\gamma$  in  $\hat{\Gamma} - \Gamma$ . Indeed, suppose that there is no such curves. That is, suppose that every non-trivial simple closed curve  $\hat{c}_i \subset \hat{c}_*$  corresponds to a conjugacy class of an element  $\gamma_i$  in  $\Gamma$ . Since  $\Gamma$  is a normal subgroup of  $\hat{\Gamma}$ , the curve  $\hat{c}_*$  corresponds to a conjugacy class of the composition of some elements in  $\{\gamma_i, \gamma_i^{-1}\}_i$ , which is in  $\Gamma$ . However, this is a contradiction. Let  $c' \subset c_*$  be a connected component of the preimage of  $\hat{c}'$ . Then  $c'$  is a simple closed curve fixed by  $f$ . If  $\gamma$  is hyperbolic, then there exists a simple closed geodesic  $c'_*$  that is homotopic to  $c'$ , and it is fixed by  $f$ . If  $\gamma$  is parabolic, then  $\hat{c}'$  surrounds a puncture  $\hat{p}$  on  $\hat{R}$ . Then  $c'$  surrounds a puncture  $p$  which is a lift of  $\hat{p}$ , and  $f$  fixes  $p$ . If  $\gamma$  is elliptic, then  $\hat{c}'$  surrounds a cone point  $\hat{p}$  on  $\hat{R}$ . In case  $c'$  is trivial, then it surrounds a point  $p$  which is a lift of  $\hat{p}$ , and  $f$  fixes  $p$ . In case  $c'$  is non-trivial, then it surrounds a cone point  $p$  which is a lift of  $\hat{p}$ , and  $f$  fixes  $p$ .  $\square$

This proposition immediately gives the following well known result.

**Corollary 1.** *Let  $R$  be a compact Riemann surface, and  $f$  a conformal automorphism of  $R$ . If  $f$  is irreducible, then  $f$  has a fixed point on  $R$ .*

Assume that  $R$  has the non-abelian fundamental group. Then the action of  $\mathrm{Aut}(R)$  is properly discontinuous (cf. [6, Theorem X.48]). Thus, if a conformal automorphism  $f$  of  $R$  fixes either a simple closed geodesic, a puncture, a point or a cone point on  $R$ , then  $f$  has the finite order. Hence, by Proposition 1,  $f$  has the finite order if and only if  $f$

fixes either a simple closed geodesic, a puncture, a point or a cone point on  $R$ . In each case, we estimate the order of  $f$  concretely in terms of the injectivity radius on  $R$ .

**Theorem 1.** (hyperbolic case) *Let  $R$  be a hyperbolic Riemann surface. Suppose that there exists a positive constant  $M$  such that the injectivity radius at any point in  $R$  is less than  $M$ . Let  $f$  be a conformal automorphism of  $R$  such that  $f(c) = c$  for a simple closed geodesic  $c$  on  $R$  whose length is  $\ell$ . Then the order  $n$  of  $f$  satisfies*

$$n < (e^{2M} - 1) \cosh(\ell/2).$$

**Theorem 2.** (parabolic case) *Let  $R$  be a hyperbolic Riemann surface. Suppose that there exists a positive constant  $M$  such that the injectivity radius at any point in  $R$  is less than  $M$ . Let  $f$  be a conformal automorphism of  $R$  such that  $f(p) = p$  for a puncture  $p$  of  $R$ . Then the order  $n$  of  $f$  satisfies*

$$n < e^{2M} - 1.$$

**Theorem 3.** (elliptic case) (i) *Let  $R$  be a hyperbolic Riemann surface, and  $f$  a conformal automorphism of  $R$  such that  $f(p) = p$  for a point  $p$  in  $R$  at which the injectivity radius is  $M > 0$ . Then the order  $n$  of  $f$  satisfies*

$$n < 2\pi \cosh M.$$

(ii) *Let  $R = \mathbb{H}/\Gamma$ , where  $\Gamma$  is a Fuchsian group which is not torsion-free. Suppose that there exists a positive constant  $M$  such that the injectivity radius at any point in  $R$  is less than  $M$ . Let  $f$  be a conformal automorphism of  $R$  such that  $f(p) = p$  for a cone point  $p$  in  $R$  which is a projection of a fixed point  $\tilde{p}$  of an elliptic element of  $\Gamma$  with order  $m > 1$ . Then the order  $n$  of  $f$  satisfies*

$$n < (e^{2M} - 1) \frac{\pi}{m} \left( \frac{1}{\sin^2 \frac{\pi}{m}} + \frac{1}{\sinh^2 M} \right)^{\frac{1}{2}}.$$

**Remark 1.** In the assumptions of Theorems 1, 2 and 3 (ii), the injectivity radius at any point in  $R$  is uniformly bounded from above. Then  $R$  must have the non-abelian fundamental group. Further, in the assumption of Theorem 3 (i), the conformal automorphism  $f$  fixes a point on  $R$  at which the injectivity radius is bounded. Thus, if the fundamental group of  $R$  is abelian, then the order of  $f$  is not greater than 2. Hence we may assume that  $R$  has the non-abelian fundamental group.

**Remark 2.** The upper bound of the order of  $f$  obtained in Theorem 2 is the limiting case of that in Theorem 1 as  $\ell \rightarrow 0$ . It is also the limiting case of that in Theorem 3 (ii) as  $m \rightarrow \infty$ .

### 3. THE COLLAR, CUSP AND CONE LEMMAS

The proofs of the theorems are based on the collar, cusp and cone lemmas (cf. [3] and [5]).

**Definition 1.** A subset  $S \subset \mathbb{H}$  is said to be *precisely invariant* under a subgroup  $\Gamma_S$  of a Fuchsian group  $\Gamma$  if  $\gamma(S) = S$  for all  $\gamma \in \Gamma_S$  and  $\gamma(S) \cap S = \emptyset$  for all  $\gamma \in \Gamma - \Gamma_S$ .

**Collar Lemma.** Let  $\Gamma$  be a Fuchsian group (which is not necessarily torsion-free) acting on  $\mathbb{H}$ , and  $L$  an axis of a hyperbolic element  $\gamma \in \Gamma$  whose translation length is less than  $\ell$ . Assume that there exists no fixed points of elements in  $\Gamma$  on  $L$  and that  $L$  is precisely invariant under the cyclic subgroup  $\langle \gamma \rangle$  generated by  $\gamma$ . Then a collar

$$C(L) = \{z \in \mathbb{H} \mid d(z, L) \leq \omega(\ell)\}$$

is precisely invariant under  $\langle \gamma \rangle$ , where  $\sinh \omega(\ell) = (2 \sinh(\ell/2))^{-1}$ . Equivalently, the boundaries  $\partial C(L)$  of  $C(L)$  and the real axis make an angle  $\theta$ , where  $\tan \theta = 2 \sinh(\ell/2)$ .

**Cusp Lemma.** Let  $\Gamma$  be a Fuchsian group (which is not necessarily torsion-free) acting on  $\mathbb{H}$ . Suppose that  $\Gamma$  contains a parabolic element  $\gamma$  with the fixed point  $\zeta$ . Then there exists a horoball  $C(\zeta)$  tangent at  $\zeta$  such that  $C(\zeta)$  is precisely invariant under the cyclic subgroup  $\langle \gamma \rangle$  generated by  $\gamma$ , and that the area of cusp neighborhood  $C(\zeta)/\langle \gamma \rangle$  is 1.

**Cone Lemma.** Let  $\Gamma$  be a Fuchsian group acting on  $\mathbb{H}$ . Suppose that a point  $p \in \mathbb{H}$  is fixed by an elliptic element  $\gamma \in \Gamma$  whose order is  $n > 2$ . Then a hyperbolic disc

$$C(p) = \{z \in \mathbb{H} \mid d(z, p) < \rho(n)\}$$

is precisely invariant under the cyclic subgroup  $\langle \gamma \rangle$  generated by  $\gamma$ , where for  $2 < n < 7$ ,  $\rho(n)$  is a constant  $\mu \approx .075$ , and for  $n \geq 7$ ,  $\cosh \rho(n) = (2 \sin(\pi/n))^{-1}$ .

## 4. PROOFS OF THE THEOREMS

In this section, we prove the theorems. First we give a proof of Theorem 1 which is based on Collar Lemma. The proof follows from the fact that there exists a wider collar of the simple closed geodesic  $c$ , as the order of a conformal automorphism  $f$  fixing  $c$  increases.

*Proof of Theorem 1:* Let  $\Gamma$  be a Fuchsian model of  $R$ , and  $\tilde{f}$  a lift of  $f$  which is a hyperbolic element in  $\mathrm{PSL}_2(\mathbb{R})$ . Note that  $\tilde{f}^n$  is a hyperbolic element in  $\Gamma$  which is corresponding to  $c$ . We consider the quotient  $\hat{R} = R/\langle f \rangle$  by the cyclic group  $\langle f \rangle$  and its Fuchsian model  $\hat{\Gamma} = \langle \Gamma, \tilde{f} \rangle$ . Then  $\hat{c} = c/\langle f \rangle$  is a simple closed geodesic on  $\hat{R}$  whose length is  $\ell/n$ . Since  $\tilde{f}$  is corresponding to  $\hat{c}$ , we may assume that  $\tilde{f}(z) = \exp(\ell/n)z$  with the axis  $L = \{iy \mid y > 0\}$ . Applying Collar Lemma for  $\hat{\Gamma}$  and  $\tilde{f}$ , we can take a collar

$$\tilde{C}(L) = \{re^{i\theta} \in \mathbb{H} \mid 0 < r, \theta_0 < \theta < \pi - \theta_0\}$$

so that it is precisely invariant under  $\langle \tilde{f} \rangle \subset \hat{\Gamma}$ , where

$$\tan \theta_0 = 2 \sinh(\ell/2n).$$

In particular,  $\gamma(\tilde{C}(L)) \cap \tilde{C}(L) = \emptyset$  for any  $\gamma \in \Gamma - \langle \tilde{f}^n \rangle$ . Then we can take a tubular neighborhood  $C(c) = \tilde{C}(L)/\langle \tilde{f}^n \rangle$  of  $c$  on  $R$  whose fundamental region is

$$A = \{re^{i\theta} \in \mathbb{H} \mid 1 < r < e^\ell, \theta_0 < \theta < \pi - \theta_0\}.$$

We may assume that  $d(c, \partial C(c)) = \omega(\ell/n) > M$ . Indeed, suppose that

$$\omega(\ell/n) = \operatorname{arcsinh} \left( \frac{1}{2 \sinh \frac{\ell}{2n}} \right) \leq M.$$

It is easily seen that

$$\frac{e^M \cosh \frac{\ell}{2}}{n} \geq \frac{\cosh \frac{\ell}{2}}{n} > \sinh \frac{\ell}{2n}$$

for  $n > 1$ ,  $\ell > 0$  and  $M > 0$ . Then

$$\frac{n}{2e^M \cosh \frac{\ell}{2}} < \frac{1}{2 \sinh \frac{\ell}{2n}} \leq \sinh M.$$

This implies that

$$\begin{aligned} n &< 2e^M \sinh M \cosh(\ell/2) \\ &= (e^{2M} - 1) \cosh(\ell/2), \end{aligned}$$

and we have nothing to prove.

We take a point  $p$  in  $C(c)$  which satisfies  $d(p, \partial C(c)) = M$ . Here  $\partial C(c)$  is a boundary curve of  $C(c)$ . From the assumption, the injectivity radius at  $p$  is less than  $M$ . That is, the length  $r_p$  of the shortest non-trivial simple closed curve  $\alpha$  passing through  $p$  is less than  $2M$ . Since  $d(p, \partial C(c)) = M$ , the curve  $\alpha$  is in  $C(c)$ . Let  $\tilde{p} = re^{i\theta} \in A$  ( $\theta_0 < \theta < \pi/2$ ) be a lift of  $p$ . Setting  $z_1(t) = re^{it}$  for  $t \geq 0$ , we have

$$M = d(\tilde{p}, \partial \tilde{C}(L)) = \int_{\theta_0}^{\theta} \frac{|z_1'(t)|}{\text{Im} z_1(t)} dt = \int_{\theta_0}^{\theta} \frac{1}{\sin t} dt \geq \int_{\theta_0}^{\theta} \frac{1}{t} dt = \log \frac{\theta}{\theta_0}.$$

Hence  $\theta \leq e^M \theta_0$ . We put  $a = e^{i\theta}$  and  $b = e^{\ell+i\theta}$ . Then  $r_p = d(a, b)$ . From Theorem 7.2.1 in [1], we have

$$\begin{aligned} \sinh \frac{1}{2} d(a, b) &= \frac{|a - b|}{2(\text{Im} a \text{Im} b)^{\frac{1}{2}}} = \frac{e^{\ell} - 1}{2e^{\frac{\ell}{2}} \sin \theta} = \frac{\sinh \frac{\ell}{2}}{\sin \theta} \geq \frac{\sinh \frac{\ell}{2}}{\theta} \\ &\geq \frac{\sinh \frac{\ell}{2}}{e^M \theta_0} = \frac{\sinh \frac{\ell}{2}}{e^M \arctan(2 \sinh \frac{\ell}{2n})} \geq \frac{\sinh \frac{\ell}{2}}{2e^M \sinh \frac{\ell}{2n}} \\ &= \frac{n \sinh \frac{\ell}{2}}{e^M \ell \sinh \frac{\ell}{2n}} \geq \frac{n \sinh \frac{\ell}{2}}{e^M \ell \sinh \ell} = \frac{n \sinh \frac{\ell}{2}}{e^M \sinh \ell} \\ &= \frac{n}{2e^M \cosh \frac{\ell}{2}}. \end{aligned}$$

For the last inequality, we used the fact that  $x(\sinh x)^{-1}$  is a monotone decreasing function for  $x > 0$ . Since  $r_p < 2M$ , this implies that

$$\begin{aligned} n &< 2e^M \sinh M \cosh(\ell/2) \\ &= (e^{2M} - 1) \cosh(\ell/2). \end{aligned}$$

□

Next, we give a proof of Theorem 2 which is based on Cusp Lemma. The idea is similar to that in Theorem 1.

*Proof of Theorem 2:* Let  $\Gamma$  be a Fuchsian model of  $R$ , and  $\tilde{f}$  a lift of  $f$  to  $\mathbb{H}$  which is a parabolic element in  $\text{PSL}_2(\mathbb{R})$ . We may assume that  $\tilde{f}(z) = z + 1$ . Note that  $\tilde{f}^n$  belongs to  $\Gamma$ . We set  $\hat{\Gamma} = \langle \Gamma, \tilde{f} \rangle$ , which is a Fuchsian model of  $\hat{R} = R/\langle f \rangle$ . Applying Cusp Lemma for  $\hat{\Gamma}$  and  $\tilde{f}$ , we can take a horoball

$$\tilde{C}(\infty) = \{z \in \mathbb{H} \mid 1 < \text{Im} z\}$$

so that it is precisely invariant under  $\langle \tilde{f} \rangle \subset \hat{\Gamma}$ . In particular, we have  $\gamma(\tilde{C}(\infty)) \cap \tilde{C}(\infty) = \emptyset$  for any  $\gamma \in \Gamma - \langle \tilde{f}^n \rangle$ . We set  $C(p) = \tilde{C}(\infty)/\langle \tilde{f}^n \rangle$ ,

whose fundamental region is

$$\{z \in \mathbb{H} \mid 0 < \operatorname{Re} z < n, 1 < \operatorname{Im} z\}.$$

We take a point  $q$  in  $C(p)$  so that  $d(q, \partial C(p)) = M$ . Here  $\partial C(p)$  is the boundary curve of  $C(p)$ . From the assumption, the injectivity radius at  $q$  is less than  $M$ . That is, the length  $r_q$  of the shortest non-trivial simple closed curve  $\alpha$  passing through  $q$  is less than  $2M$ . Since  $d(q, \partial C(p)) = M$ , the curve  $\alpha$  is in  $C(p)$ . We put  $a = e^M i$  and  $b = n + e^M i$ . Then  $r_q = d(a, b)$ . From Theorem 7.2.1 in [1], we have

$$\sinh \frac{1}{2} d(a, b) = \frac{|a - b|}{2(\operatorname{Im} a \operatorname{Im} b)^{\frac{1}{2}}} = \frac{n}{2e^M}.$$

Since  $r_q < 2M$ , this implies that

$$\begin{aligned} n &< 2e^M \sinh M \\ &= e^{2M} - 1. \end{aligned}$$

□

Finally, we prove Theorem 3. The proof is based on Cusp Lemma.

*Proof of Theorem 3 (i):* Since  $2\pi \cosh M > 6$  for  $M > 0$ , we may assume that  $n \geq 7$ . Let  $\tilde{p}$  be a lift of  $p$  to  $\mathbb{H}$ , and  $\tilde{f}$  a lift of  $f$  to  $\mathbb{H}$  fixing the point  $\tilde{p}$ , which is an elliptic element in  $\operatorname{PSL}_2(\mathbb{R})$ . Note that  $\tilde{f}^n$  is the identity. We set  $\hat{\Gamma} = \langle \Gamma, \tilde{f} \rangle$ , which is a discrete group. Applying Cone Lemma for  $\hat{\Gamma}$  and  $\tilde{f}$ , we can take a hyperbolic disc

$$\tilde{C}(\tilde{p}) = \{z \in \mathbb{H} \mid d(z, \tilde{p}) < \rho(n)\}$$

so that it is precisely invariant under  $\langle \tilde{f} \rangle \subset \hat{\Gamma}$ , where

$$\cosh \rho(n) = (2 \sin(\pi/n))^{-1}.$$

In particular,  $\gamma(\tilde{C}(\tilde{p})) \cap \tilde{C}(\tilde{p}) = \emptyset$  for any  $\gamma \in \Gamma - \{\operatorname{id}\}$ . Then there exists a hyperbolic disc  $C(p)$  centered at  $p$  with radius  $\rho(n)$ . Thus the length of any non-trivial simple closed curve passing through  $p$  is greater than  $2\rho(n)$ . On the other hand, the injectivity radius at  $p$  is  $M$  by the assumption. That is, there exists a non-trivial simple closed curve passing through  $p$  whose length is  $2M$ . Hence we have  $\rho(n) \leq M$ , and this implies that

$$\begin{aligned} n &\leq \pi \{\arcsin(2 \cosh M)^{-1}\}^{-1} \\ &< 2\pi \cosh M. \end{aligned}$$

□

*Proof of Theorem 3 (ii):* Let  $\tilde{f}$  be a lift of  $f$  to  $\mathbb{H}$ , which is an elliptic element in  $\mathrm{PSL}_2(\mathbb{R})$ . Note that the order of  $\tilde{f}$  is  $mn$ , and that  $\tilde{f}^n \in \Gamma$ . Applying Cone Lemma for  $\hat{\Gamma} = \langle \Gamma, \tilde{f} \rangle$ , we can take a hyperbolic disc

$$\tilde{C}(\tilde{p}) = \{z \in \mathbb{H} \mid d(z, \tilde{p}) < \rho(mn)\}$$

so that it is precisely invariant under  $\langle \tilde{f} \rangle \subset \hat{\Gamma}$ , where

$$\cosh \rho(mn) = (2 \sin(\pi/mn))^{-1}.$$

In particular,  $\gamma(\tilde{C}(\tilde{p})) \cap \tilde{C}(\tilde{p}) = \emptyset$  for any  $\gamma \in \Gamma - \langle \tilde{f}^n \rangle$ . Then the fundamental region of  $C(p) = \tilde{C}(\tilde{p})/\Gamma$  is conformally equivalent to

$$\{re^{i\theta} \in \mathbb{C} \mid 0 \leq r < \rho(mn), 0 < \theta < 2\pi/m\}.$$

We may assume that  $\rho(mn) > M$ . Indeed, if

$$\rho(mn) = \operatorname{arccosh}(2 \sin(\pi/mn))^{-1} \leq M,$$

then

$$\begin{aligned} n &\leq \frac{\pi}{m} (\arcsin(2 \cosh M)^{-1})^{-1} < \frac{2\pi}{m} \cosh M < \frac{2\pi e^M}{m} \\ &< \frac{2\pi e^M}{m} \left\{ \left( \frac{\sinh M}{\sin \frac{\pi}{m}} \right)^2 + 1 \right\}^{\frac{1}{2}} \\ &= (e^{2M} - 1) \frac{\pi}{m} \left( \frac{1}{\sin^2 \frac{\pi}{m}} + \frac{1}{\sinh^2 M} \right)^{\frac{1}{2}}, \end{aligned}$$

and we have nothing to prove.

We take a point  $q$  in  $C(p)$  that satisfies  $d(q, \partial C(p)) = M$ . Here  $\partial C(p)$  is the boundary curve of  $C(p)$ . From the assumption, the injectivity radius at  $q$  is less than  $M$ . That is, the length  $r_q$  of the shortest non-trivial simple closed curve  $\alpha$  passing through  $q$  is less than  $2M$ . Since  $d(q, \partial C(p)) = M$ , the curve  $\alpha$  is in  $C(p)$ . From the cosine rule of triangles (cf. [1, p.148]), we have

$$\begin{aligned} \cosh r_q &= \cosh^2(\rho(mn) - M) - \sinh^2(\rho(mn) - M) \cos(2\pi/m) \\ &= \sinh^2(\rho(mn) - M)(1 - \cos(2\pi/m)) + 1. \end{aligned}$$

Then  $r_q < 2M$  implies that

$$\begin{aligned} (1) \quad \rho(mn) &< \operatorname{arcsinh} \left( \frac{\cosh 2M - 1}{1 - \cos \frac{2\pi}{m}} \right)^{\frac{1}{2}} + M \\ &= \operatorname{arcsinh} \left( \frac{\sinh M}{\sin \frac{\pi}{m}} \right) + M \\ &=: M_1. \end{aligned}$$



To obtain (1), we used the fact that the inverse function of  $y = \sinh^2 x$  for  $x > 0$  is  $\operatorname{arcsinh} \sqrt{y}$ . Since

$$\rho(mn) = \operatorname{arccosh}(2 \sin(\pi/mn))^{-1},$$

the inequality (1) implies that

$$\begin{aligned} n &< (\pi/m) \left\{ \operatorname{arcsin}(2 \cosh M_1)^{-1} \right\}^{-1} \\ &< (2\pi/m) \cosh M_1 \\ &= (2\pi/m) \cosh \left\{ \operatorname{arcsinh} \left( \frac{\sinh M}{\sin \frac{\pi}{m}} \right) + M \right\}. \end{aligned}$$

We set

$$X = \frac{\sinh M}{\sin \frac{\pi}{m}}.$$

Using the hyperbolic cosine formula and the fact that  $\cosh(\operatorname{arcsinh} x) = (x^2 + 1)^{\frac{1}{2}}$  for  $x > 0$ , we have

$$\begin{aligned} n &< (2\pi/m) \cosh \{ \operatorname{arcsinh} X + M \} \\ &= (2\pi/m) \{ \cosh(\operatorname{arcsinh} X) \cosh M + \sinh(\operatorname{arcsinh} X) \sinh M \} \\ &< (2\pi/m) \{ \cosh(\operatorname{arcsinh} X) \cosh M + \cosh(\operatorname{arcsinh} X) \sinh M \} \\ &= (2\pi/m) e^M \cosh(\operatorname{arcsinh} X) \\ &= (2\pi/m) e^M (X^2 + 1)^{\frac{1}{2}} \\ &= \frac{2\pi e^M}{m} \left\{ \left( \frac{\sinh M}{\sin \frac{\pi}{m}} \right)^2 + 1 \right\}^{\frac{1}{2}} \\ &= (e^{2M} - 1) \frac{\pi}{m} \left( \frac{1}{\sin^2 \frac{\pi}{m}} + \frac{1}{\sinh^2 M} \right)^{\frac{1}{2}}. \end{aligned}$$

□

## 5. APPLICATION

In this section, we apply our main theorem to investigating a certain property on hyperbolic geometry on Riemann surfaces. The property we observe is as follows.

**Definition 2.** We say that a Riemann surface  $R$  satisfies the *lower bound condition* if there exists a positive constant  $\epsilon$  such that the  $\epsilon$ -thin part of  $R$  consists only of cusp neighborhoods. Further, we say that  $R$  satisfies the *upper bound condition* if there exists a positive constant  $M$  such that the injectivity radius at any point in  $R$  is less than  $M$ .

In [2], we defined the (generalized) upper bound condition, and showed the following.

**Proposition 2** ([2]). *Let  $R$  be a hyperbolic Riemann surface of analytically finite type, and  $\tilde{R}$  a normal covering surface of  $R$  which is not a universal cover. Then  $\tilde{R}$  satisfies the lower and (generalized) upper bound conditions.*

In connection with this result, we show that a Riemann surface inherits the lower and upper bound conditions from its normal covering surface.

**Proposition 3.** *Let  $R$  be a hyperbolic Riemann surface, and  $\tilde{R}$  a normal covering surface of  $R$ . If  $\tilde{R}$  satisfies the lower and upper bound conditions, then  $R$  also satisfies these conditions.*

*Proof.* It is clear that  $R$  satisfies the upper bound condition. Suppose that  $R$  does not satisfy the lower bound condition. Then  $R$  has a sequence  $\{c_n\}$  of disjoint simple closed geodesics with  $\ell_n = \ell(c_n) \rightarrow 0$  ( $n \rightarrow \infty$ ). Here  $\ell(\cdot)$  means the hyperbolic length of a curve. Let  $\tilde{c}_n \subset \tilde{R}$  be a connected component of the preimage of  $c_n$ . Since  $\tilde{R}$  satisfies the lower bound conditions, there exists a positive constant  $\epsilon$  such that  $\ell(\tilde{c}_n) > \epsilon$  for all  $n$ . We take a positive constant  $M$  so that  $\tilde{R}$  satisfies the upper bound condition for  $M$ . Assume that  $\ell(\tilde{c}_n) \leq 2M$  for infinitely many  $n$ . Then, by Theorem 1, the order of a conformal automorphism  $\tilde{f}_n$  of  $\tilde{R}$  that fixes  $\tilde{c}_n$  is less than  $N = (e^{2M} - 1) \cosh M$ . Then we have  $\ell(c_n) > \epsilon/N$ . However, this contradicts  $\ell(c_n) \rightarrow 0$  ( $n \rightarrow \infty$ ). Next, we assume that  $\ell(\tilde{c}_n) > 2M$  (including the case that  $\tilde{c}_n$  is not closed) for infinitely many  $n$ . By Collar Lemma, there exists a tubular neighborhood  $C(c_n)$  of  $c_n$  with width  $\omega(\ell_n)$ , where  $\sinh \omega(\ell_n) = (2 \sinh(\ell_n/2))^{-1}$ . From the proof of Theorem 1, there exists a (tubular) neighborhood of  $\tilde{c}_n$  with width  $\omega(\ell_n)$ . Since  $\tilde{R}$  satisfies the upper bound condition for the constant  $M$ , there exists a non-trivial simple closed curve passing through  $\tilde{p}_n \in \tilde{c}_n$  whose length is less than  $2M$ . However, since  $\ell(\tilde{c}_n) > 2M$  and since  $\omega(\ell_n) \rightarrow \infty$  as  $n \rightarrow \infty$ , we have a contradiction.  $\square$

The following example shows that, in Proposition 3, if the normal covering surface  $\tilde{R}$  of  $R$  satisfies only one of the two conditions, then  $R$  does not necessarily satisfy the conditions.

**Example 1.** Let

$$\tilde{R} = \mathbb{C} - \bigcup_{n=1}^{\infty} \bigcup_{m \in \mathbb{Z}} \left\{ \frac{m}{n} \pm n^2 \sqrt{-1} \right\},$$

and set  $R = \tilde{R}/\langle f \rangle$ , where  $f(z) = z + 1$ . The normal covering surface  $\tilde{R}$  of  $R$  satisfies the lower bound condition but does not satisfy the upper bound condition. On the other hand,  $R$  does not satisfy the lower bound condition.

*Acknowledgment:* The author would like to express her gratitude to Professors Hiroshige Shiga and Katsuhiko Matsuzaki for their valuable advices. She also thanks the referee for his useful comments.

## REFERENCES

1. A. F. Beardon, *The Geometry of Discrete Groups*, Graduate Texts in Mathematics **91**, Springer, 1983.
2. E. Fujikawa, *Limit sets and regions of discontinuity of Teichmüller modular groups*, Proc. Amer. Math. Soc., to appear.
3. N. Halpern, *A proof of the collar lemma*, Bull. London Math. Soc., **13**, 1981, 141-144.
4. W. J. Harvey, *Cyclic groups of automorphisms of a compact Riemann surface*, Quart. J. Math. **17**, 1966, 86-97.
5. J. P. Matelski, *A compactness theorem for Fuchsian groups of the second kind*, Duke Math. J. **43**, 1976, 829-840.
6. M. Tsuji, *Potential Theory in Modern Function Theory*, Chelsea, New York, 1959.

DEPARTMENT OF MATHEMATICS, TOKYO INSTITUTE OF TECHNOLOGY, OH-OKAYAMA MEGURO-KU TOKYO 152-8551, JAPAN

*E-mail address:* fujikawa@math.titech.ac.jp