



ELSEVIER

Available at  
[www.ComputerScienceWeb.com](http://www.ComputerScienceWeb.com)  
POWERED BY SCIENCE @ DIRECT®

Fuzzy Sets and Systems III (IIII) III-III

**FUZZY**  
sets and systems

[www.elsevier.com/locate/fss](http://www.elsevier.com/locate/fss)

# Lusin's theorem on fuzzy measure spaces<sup>☆</sup>

Jun Li<sup>a,\*</sup>, Masami Yasuda<sup>b</sup>

<sup>a</sup>*Department of Applied Mathematics, Southeast University, Nanjing 210096, People's Republic of China*

<sup>b</sup>*Department of Mathematics & Informatics, Faculty of Science, Chiba University, Chiba 263-8522, Japan*

Received 25 August 2002; received in revised form 5 May 2003; accepted 7 May 2003

## Abstract

In this paper, we show that weakly null-additive fuzzy measures on metric spaces possess regularity. Lusin's theorem, which is well-known in classical measure theory, is generalized to fuzzy measure space by using the regularity and weakly null-additivity. A version of Egoroff's theorem for the fuzzy measure defined on metric spaces is given. An application of Lusin's theorem to approximation in the mean of measurable function on fuzzy measure spaces is presented.

© 2003 Published by Elsevier Science B.V.

*Keywords:* Non-additive measures; Fuzzy measure; Weakly null-additivity; Regularity; Lusin's theorem; Approximation in the mean

## 1. Introduction

The well-known Lusin's theorem in classical measure theory is very important and useful for discussing the continuity and the approximation of measurable function on metric spaces [8]. Song and Li [9] investigated the regularity of null-additive fuzzy measure on metric spaces and showed Lusin's theorem on fuzzy measure space under the null-additivity condition. These improved the previous results of Wu and Ha [11]. Further discussions for the regularity of fuzzy measures were made by Pap [7], Jiang et al. [2,3], and Wu and Wu [12].

In this paper, we shall use a weaker structural characteristic of fuzzy measures—weakly null-additivity—to discuss the above-mentioned problems. Our goal is to prove the Lusin's theorem on fuzzy measure space under the weakly null-additivity condition. The paper is organized as follows. In Section 2, a necessary and sufficient condition of weakly null-additivity of fuzzy measure is presented in Lemma 1. It constitutes the essential position in our discussion here. In Section 3,

<sup>☆</sup> This work was supported by the China Scholarship Council.

\* Corresponding author. Tel.: +86-25-3792396; fax: +86-25-3792396.

E-mail address: [lijun@seu.edu.cn](mailto:lijun@seu.edu.cn) (J. Li).

1 we prove that the weakly null-additivity implies regularity for a finite fuzzy measure defined on  
 2 metric space. In Section 4, a version of Egoroff's theorem for the fuzzy measure defined on metric  
 3 spaces is given. In Section 5, by using the regularity and Egoroff's theorem we shall prove that the  
 4 well-known Lusin's theorem remains valid for those weakly null-additive fuzzy measures defined  
 5 on a metric space. These are improvements and generalizations of the earlier results of Song and  
 6 Li [9]. Lastly, as an application of Lusin's theorem, we shall describe the mean approximations of  
 7 measurable function by continuous functions, or by polynomials, or by step functions in the sense  
 of Sugeno and of Choquet integral, respectively.

## 9 2. Preliminaries

10 Throughout this paper, we suppose that  $(X, \rho)$  is a metric space, and that  $\mathcal{O}$  and  $\mathcal{C}$  are the classes  
 11 of all open and closed sets in  $(X, \rho)$ , respectively, and  $\mathcal{B}$  is Borel  $\sigma$ -algebra on  $X$ , i.e., it is the  
 12 smallest  $\sigma$ -algebra containing  $\mathcal{O}$  [1]. Unless stated otherwise all the subsets mentioned are supposed  
 13 to belong to  $\mathcal{B}$ .

A set function  $\mu: \mathcal{B} \rightarrow [0, +\infty]$  is called a *fuzzy measure*, if it satisfies the following properties:

14 (FM1)  $\mu(\emptyset) = 0$ ;

(FM2)  $A \subset B$  implies  $\mu(A) \leq \mu(B)$  (monotonicity);

15 (FM3)  $A_1 \subset A_2 \subset \dots$  implies  $\lim_{n \rightarrow \infty} \mu(A_n) = \mu(\bigcup_{n=1}^{\infty} A_n)$  (continuity from below);

16 (FM4)  $A_1 \supset A_2 \supset \dots$ , and there exists  $n_0$  with  $\mu(A_{n_0}) < +\infty$  imply

$$17 \lim_{n \rightarrow \infty} \mu(A_n) = \mu\left(\bigcap_{n=1}^{\infty} A_n\right) \quad (\text{continuity from above}).$$

18 In this paper, we always assume that  $\mu$  is a finite fuzzy measure on  $\mathcal{B}$ , i.e.,  $\mu(X) < \infty$ .

19 A fuzzy measure  $\mu$  is called *null-additive*, if  $\mu(E \cup F) = \mu(E)$  whenever  $E, F \in \mathcal{B}$  and  $\mu(F) = 0$ ;  
 20 *autocontinuous from above*, if  $\lim_{n \rightarrow +\infty} \mu(E \cup F_n) = \mu(E)$  whenever  $E \in \mathcal{B}$ ,  $\{F_n\} \subset \mathcal{B}$ , and  $\lim_{n \rightarrow +\infty}$   
 21  $\mu(F_n) = 0$  [10].

**Definition 1** (Wang and Klir [10]).  $\mu$  is called weakly null-additive, if for any  $E, F \in \mathcal{B}$ ,

$$22 \mu(E) = \mu(F) = 0 \Rightarrow \mu(E \cup F) = 0.$$

23 Obviously, the null-additivity of  $\mu$  implies weakly null-additivity. If  $\mu$  is autocontinuous from  
 24 above, then it is null-additive [10], and hence it is weakly null-additive.

**Lemma 1.**  $\mu$  is weakly null-additive if and only if for any  $\varepsilon > 0$  and any double sequence  $\{A_n^{(k)} \mid$   
 25  $n \geq 1, k \geq 1\} \subset \mathcal{B}$  satisfying  $A_n^{(k)} \searrow D_n$  ( $k \rightarrow \infty$ ),  $\mu(D_n) = 0$ ,  $n = 1, 2, \dots$ , there exists a subsequence  
 26  $\{A_n^{(k_n)}\}$  of  $\{A_n^{(k)} \mid n \geq 1, k \geq 1\}$  such that

$$27 \mu\left(\bigcup_{n=1}^{\infty} A_n^{(k_n)}\right) < \varepsilon \quad (k_1 < k_2 < \dots).$$

1 **Proof. Necessity:** Suppose  $\mu$  is weakly null-additive. Write  $D = \bigcup_{n=1}^{\infty} D_n$ , then by using the continuity  
 3 from below of  $\mu$ , we have  $\mu(D) = 0$  and  $D_n \subset D$  ( $n = 1, 2, \dots$ ). Since for any fixed  $n = 1, 2, \dots$ ,  $A_n^{(k)} \searrow$   
 $D_n$  as  $k \rightarrow \infty$ , we have

$$A_n^{(k)} \cup D \searrow D_n \cup D = D \quad (k \rightarrow \infty)$$

5 for any fixed  $n = 1, 2, \dots$ . For given  $\varepsilon > 0$ , using the continuity from above of fuzzy measures, we  
 7 have  $\lim_{k \rightarrow +\infty} \mu(A_1^{(k)} \cup D) = \mu(D) = 0$ , therefore there exists  $k_1$  such that  $\mu(A_1^{(k_1)} \cup D) < \frac{\varepsilon}{2}$ ; For this  
 $k_1$ ,

$$(A_1^{(k_1)} \cup A_2^{(k)}) \cup D \searrow (A_1^{(k_1)} \cup D_2) \cup D = A_1^{(k_1)} \cup D,$$

9 as  $k \rightarrow \infty$ . Therefore it follows, from the continuity from above of  $\mu$ , that

$$\lim_{k \rightarrow +\infty} \mu((A_1^{(k_1)} \cup A_2^{(k)}) \cup D) = \mu(A_1^{(k_1)} \cup D).$$

11 Thus there exists  $k_2 (> k_1)$ , such that

$$\mu((A_1^{(k_1)} \cup A_2^{(k_2)}) \cup D) < \frac{\varepsilon}{2}.$$

13 Generally, there exist  $k_1, k_2, \dots, k_m$ , such that

$$\mu((A_1^{(k_1)} \cup A_2^{(k_2)} \cup \dots \cup A_m^{(k_m)}) \cup D) < \frac{\varepsilon}{2}.$$

15 Hence we obtain a sequence  $\{k_n\}_{n=1}^{\infty}$  of numbers and a sequence  $\{A_n^{(k_n)}\}_{n=1}^{\infty}$  of sets. By using the  
 monotonicity and the continuity from below of  $\mu$ , we have

$$\mu \left( \bigcup_{n=1}^{+\infty} A_n^{(k_n)} \right) \leq \mu \left( \left( \bigcup_{n=1}^{+\infty} A_n^{(k_n)} \right) \cup D \right) \leq \frac{\varepsilon}{2} < \varepsilon.$$

17 *Sufficiency:* Let  $E, F \in \mathcal{B}$  and  $\mu(E) = \mu(F) = 0$ . We define a double sequence  $\{A_n^{(k)} \mid n \geq 1, k \geq 1\}$   
 19 of sets satisfying the following conditions:  $A_1^{(k)} = E$ ,  $A_2^{(k)} = F$ ,  $A_3^{(k)} = A_4^{(k)} = \dots = \emptyset$ ,  $\forall k \geq 1$  and let  
 $D_1 = E$ ,  $D_2 = F$ ,  $D_n = \emptyset$ ,  $\forall n \geq 3$ . Then for any  $\varepsilon > 0$ , by hypothesis, there exists a subsequence  $\{A_n^{(k_n)}\}$   
 21 such that  $\mu(\bigcup_{n=1}^{\infty} A_n^{(k_n)}) < \varepsilon$ , that is  $\mu(E \cup F) < \varepsilon$ . Therefore  $\mu(E \cup F) = 0$ . This shows that  $\mu$  is weakly  
 null-additive.  $\square$

23 **Remark 1.** A weakly null-additive fuzzy measure may not be null-additive. In the following, a  
 simple example indicates that the weakly null-additivity of fuzzy measure is really weaker than  
 25 null-additivity and autocontinuity from above.

**Example 1.** Let  $X = \{a, b\}$  and  $(X, \rho)$  be a metric space. Then  $\mathcal{B} = \wp(X)$ . Put

$$\mu(E) = \begin{cases} 1 & \text{if } E = X, \\ \frac{1}{2} & \text{if } E = \{b\}, \\ 0 & \text{if } E = \{a\} \text{ or } E = \emptyset. \end{cases}$$

27

1 Then  $\mu$  is a fuzzy measure with weakly null-additivity. However  $\mu$  is not null-additive and hence it  
 is not autocontinuous from above either. In fact,  $\mu(\{a\}) = 0$ , but  $\mu(\{a\} \cup \{b\}) = 1 \neq \mu(\{b\})$ .

3 **3. Regularity of fuzzy measure**

It is known that every probability measure  $P$  on a metric space is regular. Now we prove that  
 5 this property is also enjoyed by those fuzzy measures with weakly null-additivity.

**Definition 2** (Wu and Ha [11]).  $\mu$  is called regular if, for every  $A \in \mathcal{B}$  and  $\varepsilon > 0$ , there exist a closed  
 7 set  $F_\varepsilon$  and an open set  $G_\varepsilon$  of  $X$ , such that  $F_\varepsilon \subset A \subset G_\varepsilon$  and  $\mu(G_\varepsilon - F_\varepsilon) < \varepsilon$ .

**Theorem 1.** *If  $\mu$  is weakly null-additive, then  $\mu$  is regular.*

9 **Proof.** Let  $\mathcal{E}$  be the class of all set  $E \in \mathcal{B}$  such that for any  $\varepsilon > 0$ , there exist a closed set  $F_\varepsilon$  and an  
 open set  $G_\varepsilon$  satisfying

11 
$$F_\varepsilon \subset E \subset G_\varepsilon \quad \text{and} \quad \mu(G_\varepsilon - F_\varepsilon) < \varepsilon.$$

To prove the theorem, it is sufficient to show that  $\mathcal{B} \subset \mathcal{E}$ .

13 It is easy to verify that  $\emptyset \in \mathcal{E}, X \in \mathcal{E}$  and  $\mathcal{E}$  is closed under the formation of complements.

We shall now prove that  $\mathcal{E}$  is also closed under the formation of countable unions. Let  $\{E_n\} \subset \mathcal{E}$   
 15 and  $\varepsilon > 0$  be given. From the definition of  $\mathcal{E}$  and  $E_n \in \mathcal{E}$ , we know that for every  $n = 1, 2, \dots$ , there  
 exist a sequence  $\{G_n^{(k)}\}_{k=1}^\infty$  of open sets and a sequence  $\{F_n^{(k)}\}_{k=1}^\infty$  of closed sets such that

17 
$$F_n^{(k)} \subset E_n \subset G_n^{(k)} \quad \text{and} \quad \mu(G_n^{(k)} - F_n^{(k)}) < \frac{1}{k}$$

for  $k = 1, 2, \dots$ . Without loss of generality, we can assume that for fixed  $n = 1, 2, \dots$ , as  $k \rightarrow \infty$ ,  
 19  $\{G_n^{(k)}\}_{k=1}^\infty$  is decreasing and  $\{F_n^{(k)}\}_{k=1}^\infty$  is increasing. Therefore, for any fixed  $n = 1, 2, \dots$ ,  $\{G_n^{(k)} -$   
 $F_n^{(k)}\}_{k=1}^\infty$  is a decreasing sequence of sets with respect to  $k$  and as  $k \rightarrow \infty$

21 
$$G_n^{(k)} - F_n^{(k)} \searrow \bigcap_{k=1}^\infty (G_n^{(k)} - F_n^{(k)}).$$

Denote  $D_n = \bigcap_{k=1}^\infty (G_n^{(k)} - F_n^{(k)})$ , then  $G_n^{(k)} - F_n^{(k)} \searrow D_n$  as  $k \rightarrow \infty$  and noting that  $\mu(D_n) \leq \mu(G_n^{(k)} -$   
 23  $F_n^{(k)}) < \frac{1}{k}$ ,  $k = 1, 2, \dots$ , we have  $\mu(D_n) = 0$  ( $n = 1, 2, \dots$ ). Applying Lemma 1 to the double sequence  
 $\{G_n^{(k)} - F_n^{(k)}\}$  and the sequence  $\{D_n\}_{n=1}^\infty$  of sets, then for any given  $\varepsilon > 0$ , there exists a subsequence  
 25  $\{G_n^{(k_n)} - F_n^{(k_n)}\}$  of  $\{G_n^{(k)} - F_n^{(k)}\}$  such that

$$\mu \left( \bigcup_{n=1}^\infty (G_n^{(k_n)} - F_n^{(k_n)}) \right) < \varepsilon.$$

1 Since

$$\bigcup_{n=1}^{\infty} G_n^{(k_n)} - \bigcup_{n=1}^N F_n^{(k_n)} \searrow \bigcup_{n=1}^{\infty} G_n^{(k_n)} - \bigcup_{n=1}^{\infty} F_n^{(k_n)}$$

3 as  $N \rightarrow \infty$ , and noting that  $\bigcup_{n=1}^{\infty} G_n^{(k_n)} - \bigcup_{n=1}^{\infty} F_n^{(k_n)} \subset \bigcup_{n=1}^{\infty} (G_n^{(k_n)} - F_n^{(k_n)})$ , by the continuity from above and monotonicity of  $\mu$ , we have

$$5 \quad \lim_{N \rightarrow +\infty} \mu \left( \bigcup_{n=1}^{\infty} G_n^{(k_n)} - \bigcup_{n=1}^N F_n^{(k_n)} \right) = \mu \left( \bigcup_{n=1}^{\infty} G_n^{(k_n)} - \bigcup_{n=1}^{\infty} F_n^{(k_n)} \right) < \varepsilon.$$

Therefore, there exists  $N_0$  such that

$$7 \quad \mu \left( \bigcup_{n=1}^{\infty} G_n^{(k_n)} - \bigcup_{n=1}^{N_0} F_n^{(k_n)} \right) < \varepsilon.$$

Denote

$$9 \quad G_\varepsilon = \bigcup_{n=1}^{\infty} G_n^{(k_n)} \quad \text{and} \quad F_\varepsilon = \bigcup_{n=1}^{N_0} F_n^{(k_n)}$$

then  $G_\varepsilon$  is an open set,  $F_\varepsilon$  is a closed set and

$$11 \quad F_\varepsilon \subset \bigcup_{n=1}^{\infty} E_n \subset G_\varepsilon \quad \text{and} \quad \mu(G_\varepsilon - F_\varepsilon) < \varepsilon.$$

Therefore  $\bigcup_{n=1}^{\infty} E_n \in \mathcal{E}$ . Thus we proved that  $\mathcal{E}$  is a  $\sigma$ -algebra.

13 To complete the proof, it is enough to show that  $\mathcal{E}$  contains all the open sets of  $X$ . For any closed set  $F \in \mathcal{C}$ , we denote  $G_m = \{x \in X : \rho(x, F) < 1/m\}$  ( $m = 1, 2, \dots$ ), where  $\rho(x, F)$  is the distance of the set  $F$  from the point  $X$ , i.e.  $\rho(x, F) = \inf\{\rho(x, y) : y \in F\}$ , then for every  $m = 1, 2, \dots$ ,  $G_m$  is open set. Noting that  $F$  is a closed set, we know  $G_m \searrow F$  ( $m \rightarrow \infty$ ). It is follows from  $G_m - F \searrow \emptyset$  ( $m \rightarrow \infty$ ) that  $\lim_{m \rightarrow \infty} \mu(G_m - F) = 0$ . Thus  $\mathcal{C} \subset \mathcal{E}$ . Since  $\mathcal{E}$  is closed under the formation of complements, we have  $\mathcal{O} \subset \mathcal{E}$ . This shows that  $\mathcal{E}$  is a  $\sigma$ -algebra containing  $\mathcal{O}$ . Therefore  $\mathcal{B} \subset \mathcal{E}$ .  $\square$

19 **Corollary 1.** *If  $\mu$  is weakly null-additive, then for any  $E \in \mathcal{B}$ , there exist a sequence  $\{F^{(k)}\}_{k=1}^{\infty}$  of closed sets and a sequence  $\{G^{(k)}\}_{k=1}^{\infty}$  of open sets such that for every  $k = 1, 2, \dots$ ,*

$$21 \quad F^{(k)} \subset E \subset G^{(k)},$$

$$\mu(G^{(k)} - E) < \frac{1}{k} \quad \text{and} \quad \mu(E - F^{(k)}) < \frac{1}{k}.$$

23 *Note 1:* Observe that we can assume in Corollary 1 that the sequence  $\{F^{(k)}\}_{k=1}^{\infty}$  is increasing in  $k$  and the sequence  $\{G^{(k)}\}_{k=1}^{\infty}$  is decreasing in  $k$ .

1 **4. Egoroff's theorem**

3 Egoroff's theorem on fuzzy measure spaces was discussed in [4–6,10]. Now we show a version  
 5 of the Egoroff's theorem for the fuzzy measures defined on metric spaces. We assume that in this  
 7 paper all functions considered are defined on  $X$  and are real-valued measurable with respect to  $\mathcal{B}$ .  
 9 For a finite fuzzy measure  $\mu$  on  $\mathcal{B}$ , we have obtained the following result [5]:

11 **Theorem 2** (Egoroff's theorem). *If  $\{f_n\}$  converges to  $f$  almost everywhere on  $X$ , then for any  $\varepsilon > 0$   
 13 there exists  $X_\varepsilon \in \mathcal{B}$  such that  $\mu(X - X_\varepsilon) < \varepsilon$  and  $\{f_n\}_n$  converges to  $f$  uniformly on  $X_\varepsilon$ .*

The following corollary gives an alternative form of Egoroff's theorem.

15 **Corollary 2.** *If  $\{f_n\}$  converges to  $f$  almost everywhere on  $X$ , then there exists an increasing  
 17 sequence  $\{X_m\}_{m=1}^\infty \subset \mathcal{B}$  such that  $\mu(X - \bigcup_{m=1}^\infty X_m) = 0$  and  $f_n$  converges to  $f$  on  $X_m$  uniformly for  
 19 any fixed  $m = 1, 2, \dots$ .*

21 When  $\mu$  is a weakly null-additive fuzzy measure on metric space, we can obtain a slightly stronger  
 23 conclusion:

25 **Theorem 3.** *Let  $\mu$  be weakly null-additive fuzzy measure on  $\mathcal{B}$ . If  $\{f_n\}$  converges to  $f$  almost  
 27 everywhere on  $X$ , then for any  $\varepsilon > 0$  there exists a closed subset  $F_\varepsilon \in \mathcal{C}$  such that  $\mu(X - F_\varepsilon) < \varepsilon$  and  
 29  $\{f_n\}_n$  converges to  $f$  uniformly on  $F_\varepsilon$ .*

31 **Proof.** Since  $\{f_n\}$  converges to  $f$  almost everywhere on  $X$ , by using Corollary 2 there exists  
 33 an increasing sequence  $\{X_m\}_{m=1}^\infty \subset \mathcal{B}$  such that  $f_n$  converges to  $f$  on  $X_m$  uniformly for any fixed  
 35  $m = 1, 2, \dots$  and  $\mu(X - \bigcup_{m=1}^\infty X_m) = 0$ . Denote  $H = X - \bigcup_{m=1}^\infty X_m$ , then  $\mu(H) = 0$ .

37 From Corollary 1, for every fixed  $X_m$  ( $m = 1, 2, \dots$ ), there exists a sequence  $\{F_m^{(k)}\}_{k=1}^\infty$  of closed  
 39 sets satisfying  $F_m^{(k)} \subset X_m$  and  $\mu(X_m - F_m^{(k)}) < 1/k$  for any  $k = 1, 2, \dots$ . Without loss of generality, we  
 41 can assume that for fixed  $m = 1, 2, \dots$ ,  $\{X_m - F_m^{(k)}\}_{k=1}^\infty$  is decreasing (as  $k \rightarrow \infty$ ). Thus

43 
$$X_m - F_m^{(k)} \searrow \bigcap_{k=1}^\infty (X_m - F_m^{(k)})$$

45 as  $k \rightarrow \infty$ . Write  $D_m = (\bigcap_{k=1}^\infty (X_m - F_m^{(k)})) \cup H$  ( $m = 1, 2, \dots$ ), then  $(X_m - F_m^{(k)}) \cup H \searrow D_m$  as  $k \rightarrow \infty$ .  
 47 Noting that for any  $m = 1, 2, \dots$ ,  $\mu(\bigcap_{k=1}^\infty (X_m - F_m^{(k)})) = \lim_{k \rightarrow +\infty} \mu(X_m - F_m^{(k)}) = 0$ , and by the weakly  
 49 null-additivity of  $\mu$ , we get  $\mu(D_m) = \mu((\bigcap_{k=1}^\infty (X_m - F_m^{(k)})) \cup H) = 0$  ( $m = 1, 2, \dots$ ). Applying Lemma 1  
 51 to the double sequence  $\{(X_m - F_m^{(k)}) \cup H\}$  of sets and the sequence  $\{D_m\}_{m=1}^\infty$  of sets, then for any  
 53  $\varepsilon > 0$ , there exists a subsequence  $\{(X_m - F_m^{(k_m)}) \cup H\}$  of  $\{(X_m - F_m^{(k)}) \cup H\}$ , such that

55 
$$\mu \left( \bigcup_{m=1}^\infty ((X_m - F_m^{(k_m)}) \cup H) \right) < \varepsilon.$$

1 Since  $X - \bigcup_{m=1}^{\infty} F_m^{(k_m)} \subset \bigcup_{m=1}^{\infty} (X_m - F_m^{(k)}) \cup H$ , we have

$$\mu \left( X - \bigcup_{m=1}^{\infty} F_m^{(k_m)} \right) < \varepsilon.$$

3 On the other hand, from  $X - \bigcup_{m=1}^N F_m^{(k_m)} \searrow X - \bigcup_{m=1}^{\infty} F_m^{(k_m)}$  as  $N \rightarrow \infty$  and the continuity from above of  $\mu$ , we have  $\lim_{N \rightarrow +\infty} \mu(X - \bigcup_{m=1}^N F_m^{(k_m)}) = \mu(X - \bigcup_{m=1}^{\infty} F_m^{(k_m)}) < \varepsilon$ . Therefore there exists  
5  $N_0$  such that  $\mu(X - \bigcup_{m=1}^{N_0} F_m^{(k_m)}) < \varepsilon$ .

7 Denote  $F_\varepsilon = \bigcup_{m=1}^{N_0} F_m^{(k_m)}$ , then  $F_\varepsilon$  is a closed set,  $\mu(X - F_\varepsilon) < \varepsilon$  and from  $F_\varepsilon \subset \bigcup_{m=1}^{N_0} X_m$ , we know that  $\{f_n\}_n$  converges to  $f$  uniformly on  $F_\varepsilon$ .  $\square$

### 5. Lusin's theorem

9 In this section, we shall further generalize the well-known Lusin's theorem in classical measure theory to fuzzy measure space by using the results obtained in Sections 2–4.

11 **Theorem 4** (Lusin's theorem). *Let  $\mu$  be weakly null additive fuzzy measure on  $\mathcal{B}$ . If  $f$  is a real-valued measurable function on  $X$ , then, for every  $\varepsilon > 0$ , there exists a closed subset  $F_\varepsilon \in \mathcal{C}$  such  
13 that  $f$  is continuous on  $F_\varepsilon$  and  $\mu(X - F_\varepsilon) < \varepsilon$ .*

**Proof.** We prove the theorem stepwise in the following two situations.

15 (a) Suppose that  $f$  is a simple function, i.e.  $f(x) = \sum_{n=1}^s c_n \chi_{E_n}(x)$  ( $x \in X$ ), where  $\chi_{E_n}(x)$  is the characteristic function of the set  $E_n$  and  $X = \bigcup_{n=1}^s E_n$  (a disjoint finite union). For every fixed  
17  $E_n$  ( $n = 1, 2, \dots, s$ ), by Corollary 1, there exists the sequence  $\{F_n^{(k)}\}_{k=1}^{\infty}$  of closed sets such that

$$F_n^{(k)} \subset E_n \quad \text{and} \quad \mu(E_n - F_n^{(k)}) < \frac{1}{k}$$

19 for any  $k = 1, 2, \dots$ . We may assume that  $\{F_n^{(k)}\}_{k=1}^{\infty}$  is increasing in  $k$  for each fixed  $n$ , without any loss of generality.

21 For any  $\varepsilon > 0$ , applying Lemma 1 to the double sequence  $\{E_n - F_n^{(k)}\}$  ( $n = 1, 2, \dots, s, k = 1, 2, \dots$ ) of sets, there exists a subsequence  $\{E_n - F_n^{(k_n)}\}$  of  $\{E_n - F_n^{(k)}\}$  such that

$$\mu \left( \bigcup_{n=1}^s (E_n - F_n^{(k_n)}) \right) < \varepsilon.$$

Put  $F_\varepsilon = \bigcup_{n=1}^s F_n^{(k_n)}$ , then  $f$  is continuous on the closed subset  $F_\varepsilon$  of  $X$ , and

$$\mu(X - F_\varepsilon) \leq \mu \left( \bigcup_{n=1}^s E_n - \bigcup_{n=1}^s F_n^{(k_n)} \right) \leq \mu \left( \bigcup_{n=1}^s (E_n - F_n^{(k_n)}) \right) < \varepsilon.$$

25 (b) Let  $f$  be a real-valued measurable function. Then there exists a sequence  $\{\varphi_n(x)\}_{n=1}^{\infty}$  of simple functions such that  $\varphi_n \rightarrow f$  ( $n \rightarrow \infty$ ) on  $X$ . By the result obtained in (a), for each simple function  
27

1  $\varphi_n$  and every  $k = 1, 2, \dots$ , there exists closed set  $X_n^{(k)} \subset X$  such that  $\varphi_n$  is continuous on  $X_n^{(k)}$  and  
 2  $\mu(X - X_n^{(k)}) < \frac{1}{k}$  ( $k = 1, 2, \dots$ ). There is no loss of generality in assuming the sequence  $\{X_n^{(k)}\}_{k=1}^\infty$  of  
 3 closed sets is increasing with respect to  $k$  for any fixed  $n$  (otherwise, we can take  $\bigcup_{i=1}^k X_n^{(i)}$  instead  
 4 of  $X_n^{(k)}$  and noting that  $\varphi_n$  is a simple function, it remains continuous on  $\bigcup_{i=1}^k X_n^{(i)}$ ). Therefore  
 5  $X - X_n^{(k)} \searrow \bigcap_{k=1}^\infty (X - X_n^{(k)})$  as  $k \rightarrow \infty$ , and thus, we have

$$\mu \left( \bigcap_{k=1}^\infty (X - X_n^{(k)}) \right) = \lim_{n \rightarrow +\infty} \mu(X - X_n^{(k)}) = 0 \quad (n = 1, 2, \dots).$$

7 Now we consider the double sequence  $\{X - X_n^{(k)} \mid n \geq 1, k \geq 1\}$  of sets. By using Lemma 1, for  
 8 every  $m$  ( $m = 1, 2, \dots$ ), we may take a subsequence  $\{X - X_n^{(k_n^{(m)})}\}_{n=1}^\infty$  of  $\{X - X_n^{(k)} \mid n \geq 1, k \geq 1\}$  such  
 9 that

$$\mu \left( \bigcup_{n=1}^\infty (X - X_n^{(k_n^{(m)})}) \right) < \frac{1}{m},$$

11 namely,  $\mu(X - \bigcap_{n=1}^\infty X_n^{(k_n^{(m)})}) < 1/m$ . Since the double sequence  $\{X - X_n^{(k)} \mid n \geq 1, k \geq 1\}$  of sets is de-  
 12 creasing in  $k$  for fixed  $n$ , without any loss of generality, we can assume that for fixed  $n$  ( $n = 1, 2, \dots$ ),  
 13  $k_n^{(1)} < k_n^{(2)} < \dots < k_n^{(m)} \dots$ . Write  $H_m = \bigcap_{n=1}^\infty X_n^{(k_n^{(m)})}$  ( $m = 1, 2, \dots$ ), then we obtain a sequence  $\{H_m\}_{m=1}^\infty$   
 14 of closed sets satisfying  $H_1 \subset H_2 \subset \dots$  and  $\mu(X - \bigcup_{m=1}^\infty H_m) = \lim_{n \rightarrow +\infty} \mu(X - H_m) = 0$ . Noting that  
 15  $\varphi_n$  is continuous on  $X_n^{(k_n^{(m)})}$  and  $H_m \subset X_n^{(k_n^{(m)})}$  ( $n = 1, 2, \dots$ ), therefore for each  $H_m$ ,  $\varphi_n$  is continuous  
 16 on  $H_m$  for every  $n = 1, 2, \dots$ .

17 On the other hand, since  $\varphi_n \rightarrow f$  ( $n \rightarrow \infty$ ) on  $X$ , by Theorem 3, there exists an increasing sequence  
 18  $\{X_m\}_{m=1}^\infty$  of closed sets satisfying  $X - X_m \searrow X - \bigcup_{m=1}^\infty X_m$  ( $n \rightarrow +\infty$ ),  $\mu(X - \bigcup_{m=1}^\infty X_m) = 0$ , and  $\{\varphi_n\}$   
 19 converges to  $f$  uniformly on closed set  $X_m$  for every  $m = 1, 2, \dots$ .

Considering the sequence  $\{(X - H_m) \cup (X - X_m)\}_{m=1}^\infty$  of sets, then, as  $m \rightarrow +\infty$

$$21 \quad (X - H_m) \cup (X - X_m) \searrow \left( X - \bigcup_{m=1}^\infty H_m \right) \cup \left( X - \bigcup_{m=1}^\infty X_m \right).$$

By using the continuity from above and weakly null-additivity of fuzzy measures, we have

$$23 \quad \lim_{m \rightarrow +\infty} \mu((X - H_m) \cup (X - X_m)) = \mu \left( \left( X - \bigcup_{m=1}^\infty H_m \right) \cup \left( X - \bigcup_{m=1}^\infty X_m \right) \right) = 0.$$

That is,  $\lim_{m \rightarrow +\infty} \mu(X - H_m \cap X_m) = 0$ . Therefore, for given  $\varepsilon > 0$ , we can take  $m_0$  such that  $\mu(X - H_{m_0}$   
 25  $\cap X_{m_0}) < \varepsilon$ . Put  $F_\varepsilon = H_{m_0} \cap X_{m_0}$ , then  $F_\varepsilon$  is a closed set and  $\mu(X - F_\varepsilon) < \varepsilon$ . Now we show that  $f$   
 26 is continuous on  $F_\varepsilon$ . In fact,  $F_\varepsilon \subset H_{m_0}$  and  $\varphi_n$  is continuous on  $H_{m_0}$ , therefore  $\varphi_n$  is continuous on  $F_\varepsilon$   
 27 for every  $n = 1, 2, \dots$ . Noting that  $\{\varphi_n\}$  converges to  $f$  on  $F_\varepsilon$  uniformly, then  $f$  is continuous  
 on  $F_\varepsilon$ .  $\square$

29 **Remark 2.** Song and Li [9] have obtained the conclusions of Theorems 1, 3 and 4 under the  
 null-additivity condition. As shown in Example 1, weakly null-additivity is really weaker than

1 null-additivity and autocontinuity from above. Therefore, Theorems 1, 3, and 4 in this paper are  
 2 improvements of the related results in Song and Li [9] and, Wu and Ha [11].

3 **6. Applications of Lusin’s theorem**

4 Now we present some applications of Lusin’s theorem to the mean approximation of measurable  
 5 function by continuous functions, or by polynomials, or by step functions in the sense of Sugeno  
 6 and of Choquet integral, respectively.

7 Consider a nonnegative real-valued measurable function  $f$  on  $(X, \mathcal{B})$ . The *Sugeno(fuzzy) integral*  
 8 of  $f$  on  $X$  with respect to  $\mu$ , denoted by  $(S) \int f \, d\mu$ , is defined by

9 
$$(S) \int f \, d\mu = \sup_{0 \leq \alpha < +\infty} [\alpha \wedge \mu(\{x: f(x) \geq \alpha\})].$$

The *Choquet integral* of  $f$  on  $X$  with respect to  $\mu$ , denoted by  $(C) \int f \, d\mu$ , is defined by

11 
$$(C) \int f \, d\mu = \int_0^\infty \mu(\{x: f(x) > t\}) \, dt,$$

where the right side integral is Lebesgue integral.

13 We say that a measurable function sequence  $\{f_n\}_n$  converges to  $f$  in fuzzy measure  $\mu$ , and denote  
 it by  $f_n \xrightarrow{\mu} f$ , if for any  $\varepsilon > 0$ ,  $\lim_{n \rightarrow \infty} \mu(\{x: |f_n(x) - f(x)| \geq \varepsilon\}) = 0$ .

15 **Theorem 5.** *Let  $\mu$  be a weakly null-additive fuzzy measure on  $\mathcal{B}$ . If  $f$  is a real-valued measurable*  
 16 *function on  $X$ , then there exists a continuous function sequence  $\{\psi_n\}_n$  on  $X$  such that  $\psi_n \xrightarrow{\mu} f$ .*  
 17 *Furthermore, if  $|f| \leq M$ , then  $|\psi_n| \leq M$ ,  $n = 1, 2, \dots$ .*

**Proof.** For every  $n = 1, 2, \dots$ , using Theorem 4 (Lusin’s theorem), we can obtain a closed subset  
 19  $F_n$  of  $X$  such that  $f$  is continuous on  $F_n$  and  $\mu(X - F_n) < \frac{1}{n}$ . By Tietze’s extension theorem [8], for  
 every  $n = 1, 2, \dots$ , there exists continuous function  $\psi_n$  on  $X$  such that  $\psi_n(x) = f(x)$  for  $x \in F_n$ , and  
 21 if  $|f| \leq M$ , then  $|\psi_n| \leq M$ . Now we show that  $\{\psi_n\}_n$  converges to  $f$  in fuzzy measure. In fact, for  
 any  $\varepsilon > 0$ , we have  $\{x: |\psi_n(x) - f(x)| \geq \varepsilon\} \subset X - F_n$ , and therefore  $\mu(\{x: |\psi_n(x) - f(x)| \geq \varepsilon\}) \leq \mu(X -$   
 23  $F_n) < \frac{1}{n}$ ,  $n = 1, 2, \dots$ . Thus we have  $\lim_{n \rightarrow \infty} \mu(\{x: |\psi_n(x) - f(x)| \geq \varepsilon\}) = 0$ .  $\square$

25 The following result can be thought as to be the mean approximation theorem on fuzzy measure  
 spaces  $(X, \mathcal{B}, \mu)$ .

27 **Theorem 6.** *Let  $\mu$  be a weakly null-additive fuzzy measure on  $\mathcal{B}$ . If  $f$  is a real-valued measurable*  
*function on  $X$ , then there exists a continuous function sequence  $\{\psi_n\}_n$  on  $X$  such that*

$$\lim_{n \rightarrow +\infty} (S) \int |\psi_n - f| \, d\mu = 0,$$

1 Furthermore, if  $|f| \leq M$ , then  $|\psi_n| \leq M$  ( $n = 1, 2, \dots$ ) and

$$\lim_{n \rightarrow +\infty} (C) \int |\psi_n - f| d\mu = 0.$$

3 **Proof.** From Theorem 5, there exists a continuous function sequence  $\{\psi_n\}_n$  on  $X$  such that  $\psi_n \xrightarrow{\mu} f$ .  
 By using Theorem 7.4 in [10], we can directly obtain  $\lim_{n \rightarrow +\infty} (S) \int |\psi_n - f| d\mu = 0$ .

5 If  $|f| \leq M$ , then from Theorem 5,  $|\psi_n| \leq M$  ( $n = 1, 2, \dots$ ). Put

$$g_n(t) = \mu(\{x: |\psi_n(x) - f(x)| > t\}), \quad t \in [0, +\infty)$$

7 since  $\psi_n \xrightarrow{\mu} f$ , we have  $g_n(t) \xrightarrow{a.e.} 0$  on  $[0, +\infty)$  as  $n \rightarrow \infty$ . Note that  $|g_n(t)| \leq \mu(X) < \infty$ , and  $g_n(t) = 0$   
 for any  $t > 2M$  ( $n = 1, 2, \dots$ ). Applying the Bounded Convergence Theorem in Lebesgue integral  
 9 theory [8] to the function sequence  $\{g_n(t)\}_n$ , we have

$$\int_0^\infty g_n(t) dt = \int_0^{2M} g_n(t) dt \rightarrow 0 \quad (n \rightarrow \infty).$$

11 That is,  $\lim_{n \rightarrow +\infty} (C) \int |\psi_n - f| d\mu = 0$ .  $\square$

13 In the following, we discuss the mean approximation of measurable function either by polynomials  
 or by step functions on fuzzy measure space  $(R^1, \mathcal{B}, \mu)$ .

15 **Theorem 7.** Let  $\mu$  be a weakly null-additive fuzzy measure on  $\mathcal{B}$ . If  $f$  is a real-valued measurable  
 function on  $[a, b]$ , then there exists a sequence  $\{P_n\}_n$  of polynomials on  $[a, b]$  such that  $P_n \xrightarrow{\mu} f$ .  
 Furthermore, if  $|f| \leq M$ , then  $|P_n| \leq M + 1$ ,  $n = 1, 2, \dots$ .

17 **Proof.** Considering the problem on the reduced fuzzy measure space  $([a, b], [a, b] \cap \mathcal{B}, \mu)$ , then we  
 can from Theorem 5 obtain a continuous function sequence  $\{\psi_n\}_n$  on  $[a, b]$  such that  $\psi_n \xrightarrow{\mu} f$  on  
 19  $[a, b]$ . Therefore, there exists a subsequence  $\{\psi_{n_k}\}_k$  of  $\{\psi_n\}_n$ , such that

$$\mu \left( \left\{ x: |\psi_{n_k}(x) - f(x)| \geq \frac{1}{2k} \right\} \right) < \frac{1}{k},$$

21 for any  $k = 1, 2, \dots$ .

23 Since  $\psi_{n_k}$  is continuous function on  $[a, b]$  ( $k = 1, 2, \dots$ ), by using Weierstrass's theorem [8], for  
 every  $k = 1, 2, \dots$ , there exists a polynomial  $P_k$  on  $[a, b]$  such that for all  $x \in [a, b]$

$$|P_k(x) - \psi_{n_k}(x)| < \frac{1}{2k}.$$

25 Thus, for every  $k = 1, 2, \dots$ , we have

$$\left\{ x: |P_k(x) - \psi_{n_k}(x)| \geq \frac{1}{2k} \right\} = \emptyset.$$

1 Noting that

$$\left\{x: |P_k(x) - f(x)| \geq \frac{1}{k}\right\} \subset A_k \cup B_k = \left\{x: |\psi_{n_k}(x) - f(x)| \geq \frac{1}{2k}\right\},$$

3 where

$$A_k = \left\{x: |P_k(x) - \psi_{n_k}(x)| \geq \frac{1}{2k}\right\}$$

5 and

$$B_k = \left\{x: |\psi_{n_k}(x) - f(x)| \geq \frac{1}{2k}\right\},$$

7 therefore we have

$$\mu\left(\left\{x: |P_k(x) - f(x)| \geq \frac{1}{k}\right\}\right) < \frac{1}{k}.$$

9 Now we show that  $P_n \xrightarrow{\mu} f$  on  $[a, b]$ . In fact, for any given  $\varepsilon > 0$ , we take  $n_0$  such that  $1/n_0 < \varepsilon$ , then  $n \geq n_0$ ,

11 
$$\{x: |P_n(x) - f(x)| \geq \varepsilon\} \subset \left\{x: |P_n(x) - f(x)| \geq \frac{1}{n}\right\},$$

and therefore,

$$\begin{aligned} \mu(\{x: |P_n(x) - f(x)| \geq \varepsilon\}) &\leq \mu\left(\left\{x: |P_n(x) - f(x)| \geq \frac{1}{n}\right\}\right) \\ &< \frac{1}{n}, \end{aligned}$$

13 where  $n > n_0$ . This shows  $P_n \xrightarrow{\mu} f$ .

15 In the proof above, if  $|f| \leq M$ , then  $|\psi_{n_k}| \leq M$ . Since for every  $P_k$ ,  $|P_k(x) - \psi_{n_k}(x)| < \frac{1}{2k}$  for all  $x \in [a, b]$ , we have  $|P_n| \leq M + 1$ ,  $n = 1, 2, \dots$ .  $\square$

17 **Theorem 8.** Let  $\mu$  be a weakly null-additive fuzzy measure on  $\mathcal{B}$ . If  $f$  is a real-valued measurable function on  $[a, b]$ , then there exists a sequence  $\{P_n\}_n$  of polynomials on  $[a, b]$  such that

$$\lim_{n \rightarrow +\infty} (S) \int |P_n - f| d\mu = 0.$$

19 Furthermore, if  $|f| \leq M$ , then  $|P_n| \leq M + 1$  ( $n = 1, 2, \dots$ ) and

$$\lim_{n \rightarrow +\infty} (C) \int |P_n - f| d\mu = 0.$$

21 **Proof.** It is similar to the proof of Theorem 6.

1 Similarly, we can obtain the following result:

**Theorem 9.** Let  $\mu$  be a weakly null-additive fuzzy measure on  $\mathcal{B}$ . If  $f$  is a real-valued measurable function on  $[a, b]$ , then there exists a sequence  $\{s_n\}_n$  of step functions on  $[a, b]$  such that  $s_n \xrightarrow{\mu} f$  and

$$5 \quad \lim_{n \rightarrow +\infty} (S) \int |s_n - f| d\mu = 0;$$

Furthermore, if  $|f|$  is Choquet integrable, i.e.,  $(C) \int |f| d\mu < \infty$ , then  $|s_n|$  is also Choquet integrable and

$$7 \quad \lim_{n \rightarrow +\infty} (C) \int |s_n - f| d\mu = 0.$$

9 **Corollary 3.** If  $\mu$  is null-additive fuzzy measure on  $\mathcal{B}$ , then the conclusions of Theorems 5–9 hold.

## 7. Concluding remarks

11 We have proved Lusin's theorem on finite fuzzy measure space under the weakly null-additivity condition. As we have seen, the weakly null-additivity, including its a necessary and sufficient condition presented in Lemma 1, and the regularity of fuzzy measures play important roles in our discussions.

15 It should be pointed out that, in general, the weakly null-additivity is sufficient, but not necessary for Theorems 1, 4, 5 and 6 in this paper, though it is a weaker requirement, as it is weaker than null-additivity and autocontinuity.

**Example 2.** Let  $X = \{a, b\}$  and  $(X, \rho)$  be a metric space. Then  $\mathcal{B} = \wp(X)$ . Put

$$19 \quad \mu(E) = \begin{cases} 1 & \text{if } E = X, \\ 0 & \text{if } E \neq X. \end{cases}$$

Then fuzzy measure  $\mu$  is not weakly null-additive. But  $\mu$  is regular and any measurable function is continuous on  $X$ , and hence Lusin's theorem holds on  $(X, \mathcal{B}, \mu)$ .

23 We do not know whether the weakly null-additivity condition may be abandoned in our discussion. In our further research, we intend to address this issue and to investigate whether Lusin's theorem remains valid on finite fuzzy measure spaces  $(X, \mathcal{B}, \mu)$  without any additional condition as the Egoroff's theorem we have proved in [5].

## Acknowledgements

The authors are grateful to the referees and the concerned Area Editor for their valuable suggestions to revise this paper.

## 1 **References**

- 3 [1] P.R. Halmos, *Measure Theory*, Van Nostrand, New York, 1968.
- 5 [2] Q. Jiang, H. Suzuki, Fuzzy measures on metric spaces, *Fuzzy Sets and Systems* 83 (1996) 99–106.
- 7 [3] Q. Jiang, S. Wang, D. Ziou, A further investigation for fuzzy measures on metric spaces, *Fuzzy Sets and Systems* 105 (1999) 293–297.
- 9 [4] J. Li, A note on the null-additivity of the fuzzy measure, *Fuzzy Sets and Systems* 125 (2002) 269–271.
- 11 [5] J. Li, On Egoroff's theorems on fuzzy measure space, *Fuzzy Sets and Systems* 135 (2003) 367–375.
- 13 [6] J. Li, M. Yasuda, Q. Jiang, H. Suzuki, Z. Wang, G.J. Klir, Convergence of sequence of measurable functions on fuzzy measure space, *Fuzzy Sets and Systems* 87 (1997) 317–323.
- 15 [7] E. Pap, *Null-Additive Set Functions*, Kluwer, Dordrecht, 1995.
- [8] H.L. Royden, *Real Analysis*, Macmillan, New York, 1966.
- [9] J. Song, J. Li, Regularity of null-additive fuzzy measure on metric spaces, *Int. J. Gen. Systems*, to appear.
- [10] Z. Wang, G.J. Klir, *Fuzzy Measure Theory*, Plenum, New York, 1992.
- [11] C. Wu, M. Ha, On the regularity of the fuzzy measure on metric fuzzy measure spaces, *Fuzzy Sets and Systems* 66 (1994) 373–379.
- [12] J. Wu, C. Wu, Fuzzy regular measures on topological spaces, *Fuzzy Sets and Systems* 119 (2001) 529–533.